arXiv:1703.07614v1 [math.NT] 22 Mar 2017

ON THE SMALL CYCLIC TORSION OF ELLIPTIC CURVES OVER CUBIC NUMBER FIELDS.

JIAN WANG

ABSTRACT. Let E be an elliptic defined over a number field K. Then its Mordell-Weil group $E(K)$ is finitely generated: $E(K) \cong E(K)_{tor} \times \mathbb{Z}^r$. In this paper, we discuss the cyclic torsion subgroup of elliptic curves over cubic number fields. For $N = 49, 25, 32$, or $55, 40, 22, 24$, we show that $\mathbb{Z}/N\mathbb{Z}$ is not a subgroup of $E(K)_{tor}$ for any elliptic curve E over a cubic number field K.

1. INTRODUCTION

Let E be an elliptic defined over a number field K. Then its Mordell-Weil group $E(K)$ is finitely generated:

$$E(K) \cong E(K)_{tor} \times \mathbb{Z}^r$$

For a fixed elliptic E over a field K, the torsion component $E(K)_{tor}$ can be calculated due to the Nagell-Lutz-Cassels theorem [2]. However, if we consider a class of elliptic curves, it is usually difficult to list exactly all the possible group structures of $E(K)_{tor}$. The following problem is one of this kind.

Problem 1.1. *For an integer $d \geq 1$, what are the possible group structures of $E(K)_{tor}$ with $[K : \mathbb{Q}] = d$?*

For $d = 1$, i.e. $K = \mathbb{Q}$, by the work of Kubert [20] and Mazur [23], the torsion group $E(\mathbb{Q})_{tor}$ of an elliptic curve E over the rational number field is isomorphic to one of the following:

$$\begin{aligned} &\mathbb{Z}/m\mathbb{Z}, && m = 1 - 10, 12; \\ &\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2m\mathbb{Z}, && m = 1 - 4. \end{aligned}$$

For $d = 2$, by the work of Kenku-Momose [17] and Kamienny [14], the torsion group $E(K)_{tor}$ of an elliptic curve over a quadratic number field is isomorphic to one of the following:

$$\begin{aligned} &\mathbb{Z}/m\mathbb{Z}, && m = 1 - 16, 18; \\ &\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2m\mathbb{Z}, && m = 1 - 6; \\ &\mathbb{Z}/3\mathbb{Z} \times \mathbb{Z}/3m\mathbb{Z}, && m = 1 - 2; \\ &\mathbb{Z}/4\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z}. \end{aligned}$$

Date: March 23, 2017.
2010 *Mathematics Subject Classification.* 11G05,11G18.
Key words and phrases. torsion subgroup, elliptic curves, modular curves.

For $d = 3$, Parent [29, 30] showed that the prime divisors of the order of $E(K)_{tor}$ are ≤ 13. Jeon-Kim-Schweizer [12] determined all the torsion structures that appear infinitely often when we run through all elliptic curves over all cubic fields:

$$\begin{aligned} &\mathbb{Z}/m\mathbb{Z}, && m = 1 - 16, 18, 20; \\ &\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2m\mathbb{Z}, && m = 1 - 7. \end{aligned}$$

Najman [24] discovered a *sporadic* elliptic curve over a cubic field with torsion group isomorphic to $\mathbb{Z}/21\mathbb{Z}$. In view of these facts, our ultimate aim is to show that the torsion group $E(K)_{tor}$ of an elliptic curve E over a cubic number field is isomorphic to one of the following:

$$\begin{aligned} &\mathbb{Z}/m\mathbb{Z}, && m = 1 - 16, 18, 20 - 21; \\ &\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2m\mathbb{Z}, && m = 1 - 7. \end{aligned}$$

For the cyclic case, it suffices to show that $\mathbb{Z}/N\mathbb{Z}$ is not a subgroup of $E(K)_{tor}$ for any elliptic curve E over a cubic number field K when N is among the following list

$$\begin{aligned} &N = 169, 121, 49, 25, 27, 32; \\ &N = 143, 91, 65, 39, 26, 77, 55, 33, 22, 35, 63, 42, 28, 45, 30, 40, 36, 24. \end{aligned} \tag{1.1}$$

The main result of this paper is the following:

Theorem 1.2. *If $N = 49, 25, 32$, or $55, 40, 22, 24$, then $\mathbb{Z}/N\mathbb{Z}$ is not a subgroup of $E(K)_{tor}$ for any elliptic curve E over a cubic number field K.*

2. Preliminaries

Let $\mathbb{H} = \{z \in \mathbb{C} \mid \mathrm{Im} z > 0\}$ be the upper half plane. Let $\overline{\mathbb{H}} = \mathbb{H} \cup \mathbb{P}^1(\mathbb{Q})$ be the extended upper half plane by adjoining cusps $\mathbb{P}^1(\mathbb{Q}) = \mathbb{Q} \cup \{\infty\}$ to $\mathbb{H}$. Let N be a positive integer. Let $X_1(N)$ (resp. $X_0(N)$) be the modular curve defined over $\mathbb{Q}$ associated to the following congruence subgroup $\Gamma_1(N)$ (resp. $\Gamma_0(N)$) of the full modular group $\Gamma = SL_2(\mathbb{Z})/\pm 1$.

$$\Gamma_0(N) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL_2(\mathbb{Z})/\pm 1 \ \mid \ c \equiv 0 \mod N \right\}$$

$$\Gamma_1(N) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma_0(N) \ \mid \ a \equiv d \equiv 1 \mod N \right\}$$

The we have the Galois covering

$$\pi : X_1(N) \longrightarrow X_0(N)$$

with group $G = \Gamma_0(N)/\Gamma_1(N) = \left\{ \begin{pmatrix} a & 0 \\ 0 & a^{-1} \end{pmatrix} : a \in (\mathbb{Z}/N\mathbb{Z})^\times/\pm 1 \right\}$. Let Δ be a subgroup of $(\mathbb{Z}/N\mathbb{Z})^\times$ which contains ± 1. Let $X_\Delta(N)$ be the modular curve defined over $\mathbb{Q}$ associated the congruence subgroup

$$\Gamma_\Delta(N) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma_0(N) \ \mid \ a \in \Delta \mod N \right\}$$

Then all the intermediate modular curves between $X_1(N)$ and $X_0(N)$ are of the form $X_\Delta(N)$.

We denote by $Y_1(N) = X_1(N)\backslash\{cusps\}, Y_0(N) = X_0(N)\backslash\{cusps\}, Y_\Delta(N) = X_\Delta(N)\backslash\{cusps\}$ the corresponding affine curves. Denote by $J_1(N)$ (respectively, $J_0(N)$, $J_\Delta(N)$) the jacobian of $X_1(N)$ (respectively, $X_0(N)$, $X_\Delta(N)$).

The representation and behavior under the covering $\pi : X_1(N) \longrightarrow X_\Delta(N) \longrightarrow X_0(N)$ of the cusps was explained by Ogg [26]. The cusps can be regarded as pairs $\pm\binom{x}{y}$, where x, y are in $\mathbb{Z}/N\mathbb{Z}$, and relatively prime, and $\binom{x}{y}$, $\binom{-x}{-y}$ are identified. The cusps are rational in $\mathbb{Q}(\zeta_N) = \mathbb{Q}(e^{2\pi i/N})$ and cusps with the same y are conjugate. A cusp on $X_1(N)$ is an orbit $\{\pm\binom{x+by}{y}\}$, $b \in \mathbb{Z}/N\mathbb{Z}$. If $N \geq 5$, then for each $d|N$, there are $\frac{1}{2}\varphi(d)\varphi(N/d)$ cusps of $X_1(N)$ corresponding to the $\varphi(d)$ x's with $(x, d) = 1$ and the $\varphi(N/d)$ y's with $d = (y, N)$. The cusps of $X_\Delta(N)$ (and $X_0(N)$) are the orbits of G on the cusps of $X_1(N)$ with the action

$$\begin{pmatrix} a & 0 \\ 0 & a^{-1} \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} ax \\ a^{-1}y \end{pmatrix}$$

The cusps (especially of $X_0(N)$) at $d = 1$ are also called infinity cusps (denoted as ∞) and cusps at $d = N$ called zero cusps (denoted as 0).

For each divisor N' of N such that $(N', N/N') = 1$, the matrix

$$\begin{pmatrix} N'a & b \\ Nc & N'd \end{pmatrix}$$

with $a, b, c, d \in \mathbb{Z}$ and determinant N' defines a Atkin-Lehner involution $\omega_{N'}$ on $X_0(N)$. As noted in [10], it also defines an automorphism, not an involution in general, of $X_1(N)$ because it belongs to the normalizer of $\Gamma_1(N)$ in $PSL_2(\mathbb{R})$ [18]. We also denote this automorphism as $\omega_{N'}$. The Atkin-Lehner involutions are mutually commutative and satisfies the multiplication rule

$$\omega_{N_1} \circ \omega_{N_2} = \omega_{lcm(N_1,N_2)/gcd(N_1,N_2)}$$

The action of $\omega_{N'}$ on the cusps of $X_0(N)$ was given by Ogg [27, Proposition 2]. It is easy to see

$$\omega_{N'} \begin{pmatrix} 1 \\ 1 \end{pmatrix} = \begin{pmatrix} 1 \\ N' \end{pmatrix}$$

For a modular curve X, let $X^{(d)}$ be the d-th symmetric power of X, i.e. the quotient space of the d-fold product X^d by the action of the symmetric group S_d permuting the factors. Let $\overline{\mathbb{Q}}$ be the algebraic closure of $\mathbb{Q}$. Then $X^{(d)}(\overline{\mathbb{Q}})$, the set of algebraic points of $X^{(d)}$, corresponds one-to-one to the set $\{P_1 + \cdots + P_d; P_i \in X(\overline{\mathbb{Q}})\}$ of positive $\overline{\mathbb{Q}}$-rational divisors of degree d of X.

Let K be a number field of degree d over $\mathbb{Q}$. Let $x \in X(K)$. Let $x_1, \cdots, x_d$ be the images of x under the distinct embeddings $\tau_i : K \hookrightarrow \mathbb{C}, 1 \leq i \leq d$. We may view $x_1 + \cdots + x_d$ is a $\mathbb{Q}$-rational point of $X^{(d)}$. Define

$$\Phi : X^{(d)} \longrightarrow J_X$$

by $\Phi(P_1 + \cdots + P_d) = [P_1 + \cdots + P_d - d\infty]$ where J_X is the jacobian of X, and $[\ ]$ denotes the divisor class.

For a modular curve X over $\mathbb{C}$, X is called d-gonal if there exists a finite $\mathbb{C}$-morphism $\pi : X \longrightarrow \mathbb{P}^1_{\mathbb{C}}$ of degree d. The minimum d is called the *gonality* of C, which we denote as $Gon(C)$. The following lemma is a generalization of proposition 1(i) in Frey [5].

Lemma 2.1 (Frey)**.** *Assume that $Gon(X) > d$ and K is a finite extension of $\mathbb{Q}$. Then $\Phi_{|X^{(d)}(K)}$ is injective.*

Proof. Suppose otherwise $\Phi_{|X^{(d)}(K)}$ is not injective, i.e. there exists different $P_1 + \cdots + P_d$ and $Q_1 + \cdots + Q_d$ in $X^{(d)}$ such that $\Phi(P_1 + \cdots + P_d) = \Phi(Q_1 + \cdots + Q_d)$, then

$$[P_1 + \cdots + P_d - d\infty] = [Q_1 + \cdots + Q_d - d\infty] \in J_X$$

then there is a nonconstant function $f \in K(X)^*$ such that

$$\begin{aligned} \text{div}(f) &= (P_1 + \cdots + P_d - d\infty) - (Q_1 + \cdots + Q_d - d\infty) \\ &= P_1 + \cdots + P_d - Q_1 - \cdots - Q_d \end{aligned}$$

which means f has a pole divisor of degree $\leq d$. Consider the map $\pi : X \longrightarrow \mathbb{P}^1_{\mathbb{C}}$ defined by $P \longmapsto [f(P), 1]$. Then the degree of π is equal to the degree of pole divisor of f. This contradicts the assumption $Gon(X) > d$. □

We are interested in the gonality of the modular curves $X_1(N)$ over $\mathbb{C}$. Since the 1-gonal curves are precisely the curves of genus 0, then $X_1(N)$ is 1-gonal if and only if N is among the eleven values $N = 1 - 10, 12$ with genus 0. The complete lists of 2-gonal and 3-gonal ones were determined by Ishii-Momose [9] and Jeon-Kim-Schweizer [12].

Proposition 2.2 (Ishii-Momose)**.** *The modular curve $X_1(N)$ is 2-gonal if and only if N is one of the following:*

$$\begin{aligned} &N = 1 - 10, 12 \quad (g = 0); \\ &N = 11, 14, 15 \quad (g = 1); \\ &N = 13, 16, 18 \quad (g = 2). \end{aligned}$$

Proposition 2.3 (Jeon-Kim-Schweizer)**.** *The modular curve $X_1(N)$ is 3-gonal if and only if N is one of the following:*

$$\begin{aligned} &N = 1 - 10, 12 \quad (g = 0); \\ &N = 11, 14, 15 \quad (g = 1); \\ &N = 13, 16, 18 \quad (g = 2); \\ &N = 20 \qquad\quad (g = 3). \end{aligned}$$

The moduli interpretation of noncuspidal points of $X_1(N)$ are $(E, \pm P)$, where E is an elliptic curve and $P \in E$ is a point of order N. The moduli interpretation of noncuspidal points of $X_0(N)$ are (E, C), where E is an elliptic curve and $C \subset E$ is a cyclic subgroup of order N. The map $\pi : X_1(N) \longrightarrow X_0(N)$ send $(E, \pm P)$ to $(E, \langle P \rangle)$, where $\langle P \rangle$ is the cyclic subgroup generated by P.

Let p be a prime such that $p \nmid N$. Igusa's theorem [8] says that the modular curves $X_1(N)$ and $X_0(N)$ have good reduction at prime p. Moreover, reducing the modular curve is compatible with

reducing the moduli interpretation (See for example [28, Theorem 1]). And the description of the cusps is the same in characteristic p as in characteristic 0.

The description of the reduction types of elliptic curves in terms of the language of Néron models can be summarised as the Kodaira-Néron theorem [19] [25]. A complete proof of this theorem can be found in [34, IV §8 §9].

Theorem 2.4 (Kodaira-Néron)**.** *Let R be a Dedekind domain with field of fractions K, let $\mathcal{E}$ be a Néron model over R for an elliptic curve E/K, and let $\wp \subset R$ be any nonzero prime ideal with residue field k. Let $\widetilde{E}$ be the fibre over k of $\mathcal{E}$.*

(1): If E has good reduction at $\wp$, then $\widetilde{E}(k) = \widetilde{E}(k)^0$ is an elliptic curve, where $\widetilde{E}(k)^0$ denotes the connected component of the identity.

(2): If E has additive reduction at $\wp$, then $\widetilde{E}(k)^0 \cong \mathbb{G}_{a/k}$, and $\widetilde{E}(k)/\widetilde{E}(k)^0 = G$ is a finite group of order at most four.

(3): If E has multiplicative reduction at $\wp$, then there exists an extension k' of k of degree at most two so that $\widetilde{E}(k')^0 \cong \mathbb{G}_{m/k'}$ and $\widetilde{E}(k')/\widetilde{E}(k')^0 \cong \mathbb{Z}/n\mathbb{Z}$ for some positive integer n.

Let K be a number field with ring of integers $\mathcal{O}_K$, $\wp \subset \mathcal{O}_K$ a prime ideal lying above p, $k = \mathbb{F}_q = \mathcal{O}_K/\wp$ its residue field. Let E be an elliptic curve over K and $P \in E(K)$ a point of order N. Let $\widetilde{E}$ be the fibre over k of the Néron model of E, and let $\widetilde{P} \in \widetilde{E}(k)$ be the reduction of P. Suppose that $p \nmid N$. Then elementary theory of group schemes shows that $\widetilde{P}$ has order N due to the following well-known result (See for example [1, §7.3 Proposition 3]).

Proposition 2.5. *Let m be a positive integer relatively prime to $char(k)$. Then the reduction map*

$$E(K)[m] \longrightarrow \widetilde{E}(k)$$

is injective.

Let $k = \mathbb{F}_q$ be the finite field with $q = p^n$ elements. Let E/k be an elliptic curve over k. Let $|E(k)|$ be the number of points of E over k. Then Hasse's theorem states that

$$||E(k)| - q - 1| \leq 2\sqrt{q}$$

i.e.

$$(1 - \sqrt{p^n})^2 \leq |E(k)| \leq (1 + \sqrt{p^n})^2$$

Let $t = q + 1 - |E(k)|$, E is called ordinary if $(t, q) = 1$, otherwise it is called supersingular. In the range proposed by Hasse's theorem, all the ordinary t appear, while the supersingular t only appears in restricted case. This was determined by Waterhouse [35, Theorem 4.1]:

Proposition 2.6 (Waterhouse)**.** *The isogeny classes of elliptic curves over k are in one-to-one correspondence with the rational integers t having $|t| \leq 2\sqrt{q}$ and satisfying one of the following conditions:*

(1) $(t, p) = 1$;

(2) If n is even: $t = \pm 2\sqrt{q}$;

(3) If n is even and $p \not\equiv 1 \mod 3$: $t = \pm\sqrt{q}$;
(4) If n is odd and $p = 2$ or 3: $t = \pm p^{\frac{n+1}{2}}$;
(5) If either (i) n is odd or (ii) n is even and $p \not\equiv 1 \mod 4$: $t = 0$.

For the following two cases which we need in Section 4, we list here the representatives of the k-isomorphism classes of elliptic curves over k.

Case 1: $p \neq 2$ or 3, $j \neq 0, 1728$, by Silverman [33, §X Proposition 5.4], we have the following representatives of the two isomorphism classes of elliptic curves over k with $j(E_1) = j(E_2) = j$

$$E_1 : y^2 = x^3 - \frac{27j}{j-1728}x + \frac{54j}{j-1728}$$
$$E_2 : y^2 = x^3 - \frac{27j}{j-1728}\alpha^2 x + \frac{54j}{j-1728}\alpha^3$$

where $\sqrt{\alpha} \notin k^\times$. For $k = \mathbb{F}_{5^3}$, we take $\alpha = 2$.

Case 2: $p = 3$, $j \neq 0$, by Jeong [13, Theorem 3.6], we have the following representatives of the two isomorphism classes of elliptic curves over k with $j(E_1) = j(E_2) = j$

$$E_1 : y^2 = x^3 + x^2 - 1/j$$
$$E_2 : y^2 = x^3 + \alpha x^2 - \alpha^3/j$$

where $\sqrt{\alpha} \notin k^\times$. For $k = \mathbb{F}_{3^3}$, we take $\alpha = 2$.

3. Methods

The following Theorem states that the jacobian $J_1(N)$ decomposes to a direct sum of modular abelian varieties.

Theorem 3.1. [3, Theorem 6.6.6] *The jacobian $J_1(N)$ is isogenous to a direct sum of abelian varieties (over $\mathbb{Q}$) associated to equavalence classes of newforms*

$$J_1(N) \longrightarrow \bigoplus_f A_f^{m_f}$$

with $f(\tau) = \sum_{n=1}^{\infty} a_n(f) e^{2\pi i n\tau}$ newforms of divisor level.

The L-series $L(A_f, s)$ of A_f coincides, up to a finite number of Euler factors, with the product

$$\prod_\sigma L(f^\sigma, s) = \prod_\sigma \sum_{n=1}^{\infty} a_n^\sigma n^{-s}$$

where σ runs through embeddings $\sigma : K_f \hookrightarrow \mathbb{C}$ with $K_f = \mathbb{Q}(\{a_n\})$ the number field of f (See [32, §7.5]). The following proposition is a special case of Corollary 14.3 in Kato [15]:

Proposition 3.2. *Let A be an abelian variety over $\mathbb{Q}$ such that there is a surjective homomorphism $J_1(N) \longrightarrow A$ for some $N \geq 1$. If $L(A, 1) \neq 0$, then $A(\mathbb{Q})$ is finite.*

The decomposition of $J_1(N)$ and the non-vanishing of the L-series at $s = 1$ of modular abelian varieties can be calculated in Magma [21]. If $L(A_f, 1) \neq 0$ for all A_f, then we know $A_f(\mathbb{Q})$ is finite for all A_f, therefore $J_1(N)_{/\mathbb{Q}}$ is finite. For the $N \leq 65$ in the list (1.1), Table 1 is the result of calculations in Magma. The second column t is the number of non-isogenous modular abelian varieties in the decompositon $J_1(N) = \bigoplus_{i=1}^{t} A_i^{m_i}$. The third column list the dimension d_i and multiplicity m_i of each A_i (we omit m_i if $m_i = 1$). The fourth column verifies vanishing of L-series at 1 (T means $L(A_i, 1) = 0$ and F means $L(A_i, 1) \neq 0$). It is easy to see $J_1(N)(\mathbb{Q})$ is finite for the N's in Table 1 except $N = 65, 63$.

TABLE 1. Decompostion of $J_1(N)$

N	t	$d_i(m_i)$	$L(A_i, 1) = 0$
49	5	$1, 48, 6, 12, 2$	F, F, F, F, F
25	2	$8, 4$	F, F
27	2	$1, 12$	F, F
32	4	$1, 4, 8, 2(2)$	F, F, F, F
65	19	$1, 2, 2, 6, 20, 20, 8, 2(2), 8,$	$T, F, F, F, F, F, F, F, F,$
		$2, 8, 2, 8, 4, 4, 12, 6, 2, 2$	$F, F, F, F, F, F, F, F, F, F$
39	10	$1, 2, 4, 8, 2, 2(2), 4, 2, 4, 2$	$F, F, F, F, F, F, F, F, F, F$
26	5	$1, 1, 2(2), 2, 2$	F, F, F, F, F
55	10	$1, 2, 1(2), 4, 32, 8, 8, 16, 4, 4$	$F, F, F, F, F, F, F, F, F, F$
33	6	$1, 1(2), 8, 4, 4, 2$	F, F, F, F, F, F
22	2	$1(2), 4$	F, F
35	8	$1, 2, 2, 4, 4, 4, 4, 4$	F, F, F, F, F, F, F, F
63	20	$1, 2, 1(2), 6, 6, 2, 10, 4, 2(2), 2,$	$F, F, F, F, F, F, F, F, F, T$
		$10, 2, 2, 2(2), 2, 10, 2, 10, 12, 4$	$F, F, F, F, F, F, F, F, F, F$
28	4	$1(2), 4, 2, 2$	F, F, F, F
45	8	$1, 1(2), 2, 6, 16, 4, 2, 8$	F, F, F, F, F, F, F, F
30	4	$1, 1(2), 4, 2$	F, F, F, F
40	7	$1, 1(2), 4, 2(2), 8, 2, 4$	F, F, F, F, F, F, F
36	5	$1, 8, 2, 2(2), 2$	F, F, F, F, F
24	3	$1, 2, 2$	F, F, F

In the proof of Lemma 3.5, we use a specialization lemma in Appendix of Katz [16] and a theorem of Manin [22] and Drinfeld [4].

Lemma 3.3 (Specialization Lemma)**.** *Let K be a number field. Let $\wp \subset \mathcal{O}_K$ be a prime above p. Let A/K be an abelian variety. Suppose the ramification index $e_\wp(K/\mathbb{Q}) < p - 1$. Then the reduction map*

$$\Psi : A(K)_{tor} \longrightarrow A(\overline{\mathbb{F}}_p)$$

is injective.

Theorem 3.4 (Manin-Drinfeld)**.** *Let $C \subset SL_2(\mathbb{Z})/(\pm 1)$ be a congruence subgroup. $x, y \in \mathbb{P}^1(\mathbb{Q})$ and $\overline{x}, \overline{y}$ the images of x and y respectively, on $\overline{\mathbb{H}}/C$. Then the class of divisors $(\overline{x}) - (\overline{y})$ on curve $\overline{\mathbb{H}}/C$ has finite order.*

Lemma 3.5. *Suppose$N > 4$ such that Gon$(X_1(N)) > d$, $J_1(N)(\mathbb{Q})$ is finite, $p > 2$ is a prime not dividing N. Let K be a number field of degree d over $\mathbb{Q}$ and $\wp$ a prime of K over p. Let $E_{/K}$ be an elliptic curve with a K-rational point P of order N, i.e. $x = (E, \pm P) \in Y_1(N)(K)$. Then E has good reduction at $\wp$.*

Proof. Suppose E has additive reduction at $\wp$, then $\widetilde{E}(k)^0 \cong \mathbb{G}_{a/k}$ with $|\mathbb{G}_{a/k}| = p^i, i \leq d$ and $\widetilde{E}(k)/\widetilde{E}(k)^0 \cong G$ with $|G| \leq 4$. Since $\widetilde{P}$ is a k-rational point of order N in $\widetilde{E}$, then N divides $|\widetilde{E}(k)| = |\mathbb{G}_{a/k}||G|$, which is impossible under our assumption.

Suppose E has multiplicative reduction at $\wp$, i.e. x specializes to a cusp of $\widetilde{X}_1(N)$. Then $\tau_i(K)$ is also a cubic field with prime ideal $\tau_i(\wp)$ over p and residue field $k_i = k$. And $\tau_i(E)$ also has multiplicative reduction at $\tau_i(\wp)$. This means all the images $x_1, \cdots, x_d$ of x specialize to cusps of $\widetilde{X}_1(N)$. Let $c_1, \cdots, c_d$ be the cusps such that

$$x_i \otimes \overline{\mathbb{F}}_p = c_i \otimes \overline{\mathbb{F}}_p, \quad 1 \leq i \leq d$$

We know all the cusps of $X_1(N)$ are defined over $\mathbb{Q}(\zeta_N)$. Let $\wp'$ be a prime in $\mathbb{Q}(\zeta_N)$ over p. It is an elementary fact in algebraic number theory that p ramifies in $\mathbb{Q}(\zeta_N)$ if and only if $p|N$, so $e_{\wp'}(\mathbb{Q}(\zeta_N)/\mathbb{Q}) = 1$ under our assumption $p \nmid N$. So by Lemma 3.3, the specialization map

$$\Psi : J_1(N)(\mathbb{Q}(\zeta_N))_{tor} \longrightarrow J_1(N)(\overline{\mathbb{F}}_p)$$

is injective.

Since Gon$(X_1(N)) > d$, then by Lemma 2.1, the map

$$\Phi : X_1(N)^{(d)}(\mathbb{Q}(\zeta_N)) \longrightarrow J_1(N)(\mathbb{Q}(\zeta_N))$$

is injective.

We know $x_1 + \cdots + x_d$ is $\mathbb{Q}$-rational and $J_1(N)(\mathbb{Q})$ is finite, so $[x_1 + \cdots + x_d - d\infty]$ is in $J_1(N)(\mathbb{Q}(\zeta_N))_{tor}$. By Theorem 3.4, the difference of two cusps of $X_1(N)$ has finite order in $J_1(N)$. So $[c_1 + \cdots + c_d - d\infty]$ is also in $J_1(N)(\mathbb{Q}(\zeta_N))_{tor}$. Therefore $\Psi \circ \Phi(x_1 + \cdots + x_d) = \Psi \circ \Phi(c_1 + \cdots + c_d)$ implies $x_1 + \cdots + x_d = c_1 + \cdots + c_d$ since $\Psi \circ \Phi$ is injective. This is a contradiction because we assume x is a noncuspidal point.

Therefore E has good reduction at $\wp$. □

The following Proposition is used in the proofs of Lemma 3.7 and Lemma 3.8 without declaration.

Proposition 3.6. [3, Theorem 8.5.7] *Let C and C' be nonsingular projective algebraic curves over $\mathbb{Q}$ with good reduction at p and let C' have positive genus. For any morphism $h : C \longrightarrow C'$ the following diagram is commutative:*

$$\begin{array}{ccc} C & \xrightarrow{h} & C' \\ {\scriptstyle \otimes\overline{\mathbb{F}}_p}\downarrow & & \downarrow{\scriptstyle \otimes\overline{\mathbb{F}}_p} \\ \widetilde{C} & \xrightarrow{\widetilde{h}} & \widetilde{C}' \end{array}$$

Lemma 3.7. *Suppose $g(X_0(N)) > 0$. Let p be a prime not dividing N. Let K be a number field of degree d over $\mathbb{Q}$ and $\wp$ a prime of K over p. Let $E_{/K}$ be an elliptic curve with a K-rational point P of*

order N, i.e. $x = (E, \pm P) \in Y_1(N)(K)$. Suppose E has good reduction at $\wp$. Then $\omega_{N'}(\pi(x) \otimes \overline{\mathbb{F}}_p)$ also corresponds to an elliptic curve with a point of order N.

Proof. By the definitions of Atkin-Lehner involution $\omega_{N'}$ on $X_0(N)$ and the automorphism $\omega_{N'}$ on $X_1(N)$, the following diagram is commutative

$$\begin{array}{ccccc} Y_1(N) & \xrightarrow{\pi} & Y_0(N) & \xrightarrow{\otimes\overline{\mathbb{F}}_p} & \widetilde{Y}_0(N) \\ \downarrow{\scriptstyle \omega_{N'}} & & \downarrow{\scriptstyle \omega_{N'}} & & \downarrow{\scriptstyle \omega_{N'}} \\ Y_1(N) & \xrightarrow{\pi} & Y_0(N) & \xrightarrow{\otimes\overline{\mathbb{F}}_p} & \widetilde{Y}_0(N) \end{array}$$

The following diagram is also commutative

$$\begin{array}{ccc} Y_1(N) & \xrightarrow{\pi} & Y_0(N) \\ \downarrow{\scriptstyle \otimes\overline{\mathbb{F}}_p} & & \downarrow{\scriptstyle \otimes\overline{\mathbb{F}}_p} \\ \widetilde{Y}_1(N) & \xrightarrow{\pi} & \widetilde{Y}_0(N) \end{array}$$

So

$$\omega_{N'}(\pi(x) \otimes \overline{\mathbb{F}}_p) = \pi(\omega_{N'}(x)) \otimes \overline{\mathbb{F}}_p = \pi(\omega_{N'}(x) \otimes \overline{\mathbb{F}}_p)$$

Since x has good reduction, then $\omega_{N'}(x)$ also has good reduction. Now that $\omega_{N'}(x) \otimes \overline{\mathbb{F}}_p$ is the reduction of the point $\omega_{N'}(x)$ on $X_1(N)$, then it corresponds to an elliptic curve with a point of order N. Therefore $\omega_{N'}(\pi(x) \otimes \overline{\mathbb{F}}_p) = \pi(\omega_{N'}(x) \otimes \overline{\mathbb{F}}_p)$ also corresponds to an elliptic curve with a point of order N.

□

Lemma 3.8. *Suppose $g(X_0(N)) > 0$. Let p be a prime not dividing N. Let K be a cubic number field and $\wp$ a prime of K over p with residue field $k = \mathbb{F}_q = \mathbb{F}_{p^3}$. Let $E_{/K}$ be an elliptic curve with a K-rational point P of order N, i.e. $x = (E, \pm P) \in Y_1(N)(K)$. Suppose E has good reduction at $\wp$. If the Galois covering $\pi : X_1(N) \longrightarrow X_0(N)$ factors as*

$$X_1(N) \xrightarrow{\pi_1} X_\Delta(N) \xrightarrow{\pi_2} X_0(N)$$

with $g(X_\Delta(N)) = g(X_0(N)) = 1$. Then on $\widetilde{X}_0(N)$,

$$\pi(x) \otimes \overline{\mathbb{F}}_p + \phi(\pi(x) \otimes \overline{\mathbb{F}}_p) + \phi^2(\pi(x) \otimes \overline{\mathbb{F}}_p) = \pi_2(z) \otimes \overline{\mathbb{F}}_p$$

for some rational point z on $X_\Delta(N)$, where

$$\begin{aligned} \phi : \widetilde{X}_0(N) &\longrightarrow \widetilde{X}_0(N) \\ (x, y) &\longmapsto (x^p, y^p) \end{aligned}$$

is induced by the Frobenius map $\phi : \overline{\mathbb{F}}_p \longrightarrow \overline{\mathbb{F}}_p, x \longmapsto x^p$. Moreover, for any Atkin-Lehner $\omega_{N'}$,

$$\omega_{N'}(\pi(x) \otimes \overline{\mathbb{F}}_p) + \phi(\omega_{N'}(\pi(x) \otimes \overline{\mathbb{F}}_p)) + \phi^2(\omega_{N'}(\pi(x) \otimes \overline{\mathbb{F}}_p)) = \pi_2(z_{N'}) \otimes \overline{\mathbb{F}}_p$$

for some rational point $z_{N'}$ on $X_\Delta(N)$.

Proof. Consider the Galois closure L of K. Then either $\mathrm{Gal}(L/\mathbb{Q}) = \mathbb{Z}/3\mathbb{Z}$ or $\mathrm{Gal}(L/\mathbb{Q}) = S_3$. We claim that there is an element $\sigma \in \mathrm{Gal}(L/\mathbb{Q})$ of order 3, such that (after a necessary rearrangement) the embeddings $\tau_i : K \hookrightarrow \mathbb{C}, 1 \leq i \leq 3$ satisfy

$$\tau_i = \sigma^i|_K$$

In fact, if $\mathrm{Gal}(L/\mathbb{Q}) = \mathbb{Z}/3\mathbb{Z}$, then $L = K$, i.e. $K/\mathbb{Q}$ is a Galois extension. So $\mathrm{Gal}(K/\mathbb{Q}) = \{\tau_1, \tau_2, \tau_3\}$. Let σ be a generator of $\mathrm{Gal}(K/\mathbb{Q}) = \mathbb{Z}/3\mathbb{Z}$. Then, after a necessary rearrangement of τ_1, τ_2 and τ_3, we have $\tau_i = \sigma^i$.

Otherwise, if $\mathrm{Gal}(L/\mathbb{Q}) = S_3$, then L/K is a quadratic extension. There is an element $\sigma_2 \in \mathrm{Gal}(L/\mathbb{Q})$ of order 2 such that $\mathrm{Gal}(L/K) = \langle\sigma_2\rangle$. Let $\sigma_3 \in \mathrm{Gal}(L/\mathbb{Q})$ be an element of order 3. Then $\mathrm{Gal}(L/\mathbb{Q}) = \langle\sigma_2, \sigma_3\rangle$. On the other hand, each τ_i extends to two embedding $\tau_{i1}, \tau_{i2} : L \hookrightarrow \mathbb{Q}$ and $\mathrm{Gal}(L/\mathbb{Q}) = \{\tau_{11}, \tau_{12}, \tau_{21}, \tau_{22}, \tau_{31}, \tau_{32}\}$. Without loss of generality, suppose τ_3 is the identity embedding, then $\{\tau_{31}, \tau_{32}\} = \{id, \sigma_2\}$, and after a necessary rearrangement of τ_1 and τ_2, $\{\tau_{11}, \tau_{12}\} = \{\sigma_3, \sigma_3\sigma_2\}, \{\tau_{21}, \tau_{22}\} = \{\sigma_3^2, \sigma_3^2\sigma_2\}$. Let $\sigma = \sigma_3$. Then $\tau_i = \sigma^i|_K$.

The projection of the rational divisor class $[x_1+x_2+x_3-d\infty] \in J_1(N)$ on $J_\Delta(N)$ is also rational. i.e.

$$\begin{aligned}\pi_{1*}(\big[\sigma(x) + \sigma^2(x) + \sigma^3(x) - d\infty\big]) &= \pi_{1*}([\tau_1(x) + \tau_2(x) + \tau_3(x) - d\infty]) \\ &= \pi_{1*}([x_1 + x_2 + x_3 - d\infty]) \in J_\Delta(N)(\mathbb{Q})\end{aligned}$$

Since the genus of $X_\Delta(N)$ is 1, i.e. $X_\Delta(N)$ is an elliptic curve, then there is a canonical isomorphism

$$\begin{aligned}\iota : J_\Delta(N) &\longrightarrow X_\Delta(N) \\ \left[\sum_i n_i x_i\right] &\longmapsto \sum_i n_i x_i\end{aligned}$$

where the right $\sum_i$ is the group law on the elliptic curve $X_\Delta(N)$. Since the Galois action is compatible with the projection, i.e. the following diagram is commutive

$$\begin{array}{ccc} J_1(N) & \xrightarrow{\pi_{1*}} & J_\Delta(N) \\ \downarrow{\scriptstyle\sigma} & & \downarrow{\scriptstyle\sigma} \\ J_1(N) & \xrightarrow{\pi_{1*}} & J_\Delta(N) \end{array}$$

then

$$\begin{aligned}&\iota \circ \pi_{1*}(\big[\sigma(x) + \sigma^2(x) + \sigma^3(x) - d\infty\big]) \\ =&\iota(\big[\sigma(\pi_1(x)) + \sigma^2(\pi_1(x)) + \sigma^3(\pi_1(x)) - d\infty\big]) \\ =&\sigma(\pi_1(x)) + \sigma^2(\pi_1(x)) + \sigma^3(\pi_1(x)) \in X_\Delta(N)(\mathbb{Q})\end{aligned}$$

i.e. there is a rational point z on $X_\Delta(N)$ such that

$$\sigma(\pi_1(x)) + \sigma^2(\pi_1(x)) + \sigma^3(\pi_1(x)) = z$$

Let $\wp'$ be a prime of L over $\wp$ with residue field $k' = \mathcal{O}_L/\wp'$ (which is an extension of k). It is known in algebraic number theory that the Frobenius $\phi \in \mathrm{Gal}(k'/\mathbb{F}_p)$ is the reduction from a Frobenius element $Frob_{\wp'}$ in $\mathrm{Gal}(L/\mathbb{Q})$. It is easy to see $k' = k$ since the highest order of an element

in Gal$(L/\mathbb{Q})$ is 3. Now, we have an element $\sigma'(:= Frob_{\wp'}) \in$ Gal$(L/\mathbb{Q})$ of order 3 such that the following diagram is commutative:

$$\begin{array}{ccccc} X_\Delta(N) & \xrightarrow{\pi_2} & X_0(N) & \xrightarrow{\otimes\overline{\mathbb{F}}_p} & \widetilde{X}_0(N) \\ \downarrow{\scriptstyle \sigma'} & & \downarrow{\scriptstyle \sigma'} & & \downarrow{\scriptstyle \phi} \\ X_\Delta(N) & \xrightarrow{\pi_2} & X_0(N) & \xrightarrow{\otimes\overline{\mathbb{F}}_p} & \widetilde{X}_0(N) \end{array}$$

It is easy to know either $\sigma' = \sigma$ or $\sigma' = \sigma^2$. In both cases, we always have

$$id + \sigma' + \sigma'^2 = \sigma + \sigma^2 + \sigma^3$$

Since both $X_\Delta(N)$ and $X_0(N)$ are elliptic curves, they are equipped with the group law. We can always choose an infinity cusp ∞_Δ on $X_\Delta(N)$ as the zero point O_Δ and the unique infinity cusp ∞_1 on $X_0(N)$ as the zero point O_1. Since the projection $\pi_2 : X_\Delta(N) \longrightarrow X_0(N)$ maps ∞_Δ to ∞_1, then it is an isogeny, thus preserves the group law. And our assumption $p \nmid N$ guarantees $X_0(N)$ have good reduction at $\wp$. Then the group law is also preserved during reduction. In other words, the following diagram is commutative:

$$\begin{array}{ccccc} X_\Delta(N) & \xrightarrow{\pi_2} & X_0(N) & \xrightarrow{\otimes\overline{\mathbb{F}}_p} & \widetilde{X}_0(N) \\ \downarrow{\scriptstyle +} & & \downarrow{\scriptstyle +} & & \downarrow{\scriptstyle +} \\ X_\Delta(N) & \xrightarrow{\pi_2} & X_0(N) & \xrightarrow{\otimes\overline{\mathbb{F}}_p} & \widetilde{X}_0(N) \end{array}$$

Therefore

$$\begin{aligned} &\pi(x) \otimes \overline{\mathbb{F}}_p + \phi(\pi(x) \otimes \overline{\mathbb{F}}_p) + \phi^2(\pi(x) \otimes \overline{\mathbb{F}}_p) \\ =&(\pi_2 \circ \pi_1(x)) \otimes \overline{\mathbb{F}}_p + \phi((\pi_2 \circ \pi_1(x)) \otimes \overline{\mathbb{F}}_p) + \phi^2((\pi_2 \circ \pi_1(x)) \otimes \overline{\mathbb{F}}_p) \\ =&\pi_2(\pi_1(x)) \otimes \overline{\mathbb{F}}_p + \pi_2(\sigma'(\pi_1(x))) \otimes \overline{\mathbb{F}}_p + \pi_2(\sigma'^2(\pi_1(x))) \otimes \overline{\mathbb{F}}_p \\ =&(\pi_2(\pi_1(x)) + \pi_2(\sigma'(\pi_1(x))) + \pi_2(\sigma'^2(\pi_1(x)))) \otimes \overline{\mathbb{F}}_p \\ =&\pi_2(\pi_1(x) + \sigma'(\pi_1(x)) + \sigma'^2(\pi_1(x))) \otimes \overline{\mathbb{F}}_p \\ =&\pi_2(\sigma(\pi_1(x)) + \sigma^2(\pi_1(x)) + \sigma^3(\pi_1(x))) \otimes \overline{\mathbb{F}}_p \\ =&\pi_2(z) \otimes \overline{\mathbb{F}}_p \end{aligned}$$

as desired. Moreover, for any Atkin-Lehner involution $\omega_{N'}$, we have

$$\omega_{N'}(\pi(x) \otimes \overline{\mathbb{F}}_p) = \pi(\omega_{N'}(x)) \otimes \overline{\mathbb{F}}_p$$

because of the following commutative diagram:

$$\begin{array}{ccccc} Y_1(N) & \xrightarrow{\pi} & Y_0(N) & \xrightarrow{\otimes\overline{\mathbb{F}}_p} & \widetilde{Y}_0(N) \\ \downarrow{\scriptstyle \omega_{N'}} & & \downarrow{\scriptstyle \omega_{N'}} & & \downarrow{\scriptstyle \omega_{N'}} \\ Y_1(N) & \xrightarrow{\pi} & Y_0(N) & \xrightarrow{\otimes\overline{\mathbb{F}}_p} & \widetilde{Y}_0(N) \end{array}$$

Therefore

$$\begin{aligned}&\omega_{N'}(\pi(x)\otimes\overline{\mathbb{F}}_p)+\phi(\omega_{N'}(\pi(x)\otimes\overline{\mathbb{F}}_p))+\phi^2(\omega_{N'}(\pi(x)\otimes\overline{\mathbb{F}}_p))\\ =&\pi(\omega_{N'}(x))\otimes\overline{\mathbb{F}}_p+\phi(\pi(\omega_{N'}(x))\otimes\overline{\mathbb{F}}_p)+\phi^2(\pi(\omega_{N'}(x))\otimes\overline{\mathbb{F}}_p)\\ =&\pi_2(z_{N'})\otimes\overline{\mathbb{F}}_p\end{aligned}$$

for some rational point $z_{N'}$ on $X_\Delta(N)$. □

4. Proof of Theorem 1.2

4.1. $N=49,55,40$**.** As is seen in Table 1, $J_1(N)(\mathbb{Q})$ is finite. By Proposition 2.2 and 2.3, we know $Gon(X_1(N))>3$. Let K be a cubic field and $\wp$ a prime of K over 3. Suppose $x=(E,\pm P)\in Y_1(N)(K)$. Therefore by Lemma 3.5, E has good reduction at $\wp$. By Proposition 2.5, the reduction $\widetilde{P}$ of P is a k-rational point of order N in the elliptic curve $\widetilde{E}$ over $k=\mathcal{O}_K/\wp$.

But $\widetilde{E}(k)$ can not have a point of order N since $N>(1+\sqrt{3^3})^2\approx 38.4$. This is a contradiction. So $\mathbb{Z}/N\mathbb{Z}$ is not a subgroup of $E(K)_{tor}$.

4.2. $N=25,22$**.** As is seen in Table 1, $J_1(N)(\mathbb{Q})$ is finite. By Proposition 2.2 and 2.3, we know $Gon(X_1(N))>3$. Let K be a cubic field and $\wp$ a prime of K over 3. Suppose $x=(E,\pm P)\in Y_1(N)(K)$. Therefore by Lemma 3.5, E has good reduction at $\wp$. By Proposition 2.5, the reduction $\widetilde{P}$ of P is a k-rational point of order N in the elliptic curve $\widetilde{E}$ over $k=\mathcal{O}_K/\wp$.

If $k=\mathbb{F}_3$ or $\mathbb{F}_{3^2}$, then $\widetilde{E}(k)$ can not have a point of order N since $N>(1+\sqrt{3^2})^2$. If $k=\mathbb{F}_{3^3}$, suppose $\widetilde{E}(k)$ has a point of order N, then $\widetilde{E}(k)\cong\mathbb{Z}/N\mathbb{Z}$ since $Nm>(1+\sqrt{3^3})^2$ for any $m>1$. But by Theorem 2.6, $|\widetilde{E}(k)|\neq N$ ($t=3$ for $N=25$, $t=6$ for $N=22$). This is a contradiction. So $\mathbb{Z}/N\mathbb{Z}$ is not a subgroup of $E(K)_{tor}$.

4.3. $N=32$**.** As is seen in Table 1, $J_1(N)(\mathbb{Q})$ is finite. By Proposition 2.2 and 2.3, we know $Gon(X_1(N))>3$. Let K be a cubic field and $\wp$ a prime of K over 3. Suppose $x=(E,\pm P)\in Y_1(N)(K)$. Therefore by Lemma 3.5, E has good reduction at $\wp$. By Proposition 2.5, the reduction $\widetilde{P}$ of P is a k-rational point of order N in the elliptic curve $\widetilde{E}$ over $k=\mathcal{O}_K/\wp$.

The following lemma shows that $\widetilde{E}(k)$ can not have a point of order 32. This contradiction implies that $\mathbb{Z}/32\mathbb{Z}$ is not a subgroup of $E(K)_{tor}$. The calculations on the finite field $\mathbb{F}_{3^3}$ in this Lemma is done in Sage [31].

Lemma 4.1. *$\widetilde{E}(k)$ can not have a point of order 32*

Proof. For the modular curve $X_0(32)$, Fricke [6] calculated the defining equation and the explicit formula of j-invariant and the Atkin-Lehner ω_{32} on this equation. These data can also be found in Furumoto-Hasegawa [7]. The equation is

$$y^2=x^3+6x^2+16x+16 \tag{4.1}$$

with the formula of ω_{32}

$$\omega_{32}(P)=-P\pm(0,4)$$

and the formula of j-invariant

$$j = \frac{256((x(x+4)/2)^4 + 8(x(x+4)/2)^3 + 20(x(x+4)/2)^2 + 16(x(x+4)/2) + 1)^3}{(x(x+4)/2)((x(x+4)/2) + 4)((x(x+4)/2) + 2)^2}$$

It can be calculated in Sage [31] that the set of rational torsion points of this equation is

$$X_0(32)(\mathbb{Q})_{tor} = \{(\infty, \infty), (0, 4), (-2, 0), (0, -4)\} \quad \cong \mathbb{Z}/4\mathbb{Z}$$

This is exactly all the rational points of this equation since $X_0(32)(\mathbb{Q}) \cong J_0(32)(\mathbb{Q})$ is finite.

The map $\pi : X_1(32) \longrightarrow X_0(32)$ factors through $X_\Delta(32)$, where $\Delta = \{\pm 1, \pm 7, \pm 9, \pm 15\}$, with $g(X_\Delta(32)) = 1$ (See for example [11]). The cusps of $X_\Delta(32)$ over the 0 cusp of $X_0(32)$ are quadratic. This is because the eight cusps of $X_1(32)$ over 0 are

$$\pm\begin{pmatrix} 1 \\ 0 \end{pmatrix}, \pm\begin{pmatrix} 3 \\ 0 \end{pmatrix}, \pm\begin{pmatrix} 5 \\ 0 \end{pmatrix}, \pm\begin{pmatrix} 7 \\ 0 \end{pmatrix}, \pm\begin{pmatrix} 9 \\ 0 \end{pmatrix}, \pm\begin{pmatrix} 11 \\ 0 \end{pmatrix}, \pm\begin{pmatrix} 13 \\ 0 \end{pmatrix}, \pm\begin{pmatrix} 15 \\ 0 \end{pmatrix}$$

They have two orbits under the action of Δ:

$$\left\{\pm\begin{pmatrix} 1 \\ 0 \end{pmatrix}, \pm\begin{pmatrix} 7 \\ 0 \end{pmatrix}, \pm\begin{pmatrix} 9 \\ 0 \end{pmatrix}, \pm\begin{pmatrix} 15 \\ 0 \end{pmatrix}\right\}$$
$$\left\{\pm\begin{pmatrix} 3 \\ 0 \end{pmatrix}, \pm\begin{pmatrix} 5 \\ 0 \end{pmatrix}, \pm\begin{pmatrix} 11 \\ 0 \end{pmatrix}, \pm\begin{pmatrix} 13 \\ 0 \end{pmatrix}\right\}$$

So $X_\Delta(32)$ has two cusps $\pm\binom{1}{0}$, $\pm\binom{3}{0}$ over 0. They are conjugates since they have the same y in the representation $\pm\binom{x}{y}$.

Now if $\omega_{32}(P) = -P + (0, 4)$, then $(0, 4) = \omega_{32}(\infty) = 0$, which means $(0, 4)$ comes from quadratic cusps of $X_0(32)$ under $\pi_2 : X_\Delta(32) \longrightarrow X_0(32)$. If $\omega_{32}(P) = -P - (0, 4) = -P + (0, -4)$, then $(0, -4) = \omega_{32}(\infty) = 0$, which means $(0, -4)$ comes from quadratic cusps of $X_0(32)$ under $\pi_2 : X_\Delta(32) \longrightarrow X_0(32)$.

If $k = \mathbb{F}_3$ or $\mathbb{F}_{3^2}$, then $\widetilde{E}(k)$ can not have a point of order 32 since $32 > (1 + \sqrt{3^2})^2$. If $k = \mathbb{F}_{3^3}$, let $[m, n, l]$ denote the element $ma^2 + na + l \in \mathbb{F}_{3^3} = \mathbb{F}_3[a]$. Table 2 list the coordinates and j-invariant of the points on the reduction of the equation 4.1 modulo 3 which are rational over $\mathbb{F}_{3^3}$. Denote $\varphi = id + \phi + \phi^2$. The last two columns shows $\varphi(X, Y)$.

Table 3 shows the group structures of elliptic curves over $\mathbb{F}_{3^3}$ with the given j-invariants that appear in Table 2. It is easy to see that only the point P with j-invariants marked by $(*)$ may correspond to an elliptic curve with a point of order 32.

Denote $\omega(P) = -P + (0, 4) \equiv -P + (0, 1) \mod 3$ and $\omega'(P) = -P - (0, 4) = -P + (0, -4) \equiv -P + (0, 2) \mod 3$. By using the additive law on $X_0(32)$, both $\omega(P)$ and $\omega'(P)$ are calculated as listed in Table 4.

Consider the points $P = (X, Y)$ numbered by $1, \cdots, 12$ with $j(P)$ among those marked by $(*)$. Now, if $\omega_{32} = \omega$, then only the even numbered points 2,4,6,8,10,12 satisfy that $j(\omega_{32}(P))$ is among those marked by $(*)$ in Table 3, which is required by Lemma 3.7. But they have

$$\varphi(P) = (0, 1) \qquad (2, 4, 6)$$
$$\varphi(\omega_{32}(P)) = (0, 1) \qquad (8, 10, 12)$$

TABLE 2. Points on $\widetilde{X}_0(32)(\mathbb{F}_{3^3})$

X	Y_1	Y_2	j	$\varphi(X,Y_1)$	$\varphi(X,Y_2)$
∞	∞	∞	$-$		
0	2	1	$-$		
1	0	0	$-$		
$[0,1,0]$	$[2,1,1]$	$[1,2,2]$	$[2,1,2](*)$	$(0,2)$	$(0,1)$
$[0,1,1]$	$[2,2,1]$	$[1,1,2]$	$[2,2,2](*)$	$(0,2)$	$(0,1)$
$[0,1,2]$	$[2,0,2]$	$[1,0,1]$	$[2,0,0](*)$	$(0,2)$	$(0,1)$
$[1,0,1]$	$[1,1,0]$	$[2,2,0]$	$[2,1,2](*)$	$(1,0)$	$(1,0)$
$[1,1,2]$	$[1,2,0]$	$[2,1,0]$	$[2,0,0](*)$	$(1,0)$	$(1,0)$
$[1,2,2]$	$[1,0,2]$	$[2,0,1]$	$[2,2,2](*)$	$(1,0)$	$(1,0)$
$[2,0,0]$	$[1,0,0]$	$[2,0,0]$	$[2,2,1]$	(∞,∞)	(∞,∞)
$[2,0,2]$	$[0,2,2]$	$[0,1,1]$	$[2,1,1]$	$(0,2)$	$(0,1)$
$[2,1,1]$	$[0,2,1]$	$[0,1,2]$	$[2,2,1]$	$(0,2)$	$(0,1)$
$[2,1,2]$	$[1,2,1]$	$[2,1,2]$	$[2,0,2]$	(∞,∞)	(∞,∞)
$[2,2,1]$	$[0,2,0]$	$[0,1,0]$	$[2,0,2]$	$(0,2)$	$(0,1)$
$[2,2,2]$	$[1,1,1]$	$[2,2,2]$	$[2,1,1]$	(∞,∞)	(∞,∞)

TABLE 3. Group structures of elliptic curves over $\mathbb{F}_{3^3}$

j	$E_1(\mathbb{F}_{3^3})$	$E_2(\mathbb{F}_{3^3})$	j	$E_1(\mathbb{F}_{3^3})$	$E_2(\mathbb{F}_{3^3})$
$[2,0,0](*)$	$\mathbb{Z}/24$	$\mathbb{Z}/32$	$[2,0,2]$	$\mathbb{Z}/2\times\mathbb{Z}/12$	$\mathbb{Z}/2\times\mathbb{Z}/16$
$[2,1,1]$	$\mathbb{Z}/2\times\mathbb{Z}/12$	$\mathbb{Z}/2\times\mathbb{Z}/16$	$[2,1,2](*)$	$\mathbb{Z}/24$	$\mathbb{Z}/32$
$[2,2,1]$	$\mathbb{Z}/2\times\mathbb{Z}/12$	$\mathbb{Z}/2\times\mathbb{Z}/16$	$[2,2,2](*)$	$\mathbb{Z}/24$	$\mathbb{Z}/32$

which is not allowed by Lemma 3.8.

Otherwise, if $\omega_{32}=\omega'$, then only the odd numbered points 1,3,5,7,9,11 satisfy that $j(\omega_{32}(P))$ is among those marked by $(*)$ in Table 3, which is required by Lemma 3.7. But they have

$$\varphi(P)=(0,2) \qquad (1,3,5)$$
$$\varphi(\omega_{32}(P))=(0,2) \qquad (7,9,11)$$

which is not allowed by Lemma 3.8.

Actually, the information in Table 4 can be represented in Figure 1 and Figure 2. The black dot means either the point has unmarked j-invariant or the point has $\varphi(P)$ not allowed by Lemma 3.8. The white dot point is satisfactory, but each of them is connected to a black dot point by ω_{32}, which is not allowed by Lemma 3.7 or Lemma 3.8.

□

4.4. $N=24$**.** As is seen in Table 1, $J_1(N)(\mathbb{Q})$ is finite. By Proposition 2.2 and 2.3, we know $Gon(X_1(N))>3$. Let K be a cubic field and $\wp$ a prime of K over 5. We can always choose $\wp$ such that the residue field $k=\mathcal{O}_K/\wp$ has degree 1 or 3 over $\mathbb{F}_p$. In fact, the decomposition of p in $\mathcal{O}_K$

TABLE 4. Atkin-Lehner involution on $\widetilde{X}_0(32)(\mathbb{F}_{3^3})$

P	$X(P)$	$Y(P)$	$X(\omega(P))$	$Y(\omega(P))$	$X(\omega'(P))$	$Y(\omega'(P))$
1	$[0,1,0]$	$[2,1,1]$	$[2,2,2]$	$[2,2,2]$	$[1,0,1]$	$[1,1,0]$
2	$[0,1,0]$	$[1,2,2]$	$[1,0,1]$	$[2,2,0]$	$[2,2,2]$	$[1,1,1]$
3	$[0,1,1]$	$[2,2,1]$	$[2,0,0]$	$[2,0,0]$	$[1,2,2]$	$[1,0,2]$
4	$[0,1,1]$	$[1,1,2]$	$[1,2,2]$	$[2,0,1]$	$[2,0,0]$	$[1,0,0]$
5	$[0,1,2]$	$[2,0,2]$	$[2,1,2]$	$[2,1,2]$	$[1,1,2]$	$[1,2,0]$
6	$[0,1,2]$	$[1,0,1]$	$[1,1,2]$	$[2,1,0]$	$[2,1,2]$	$[1,2,1]$
7	$[1,0,1]$	$[1,1,0]$	$[2,2,1]$	$[0,1,0]$	$[0,1,0]$	$[2,1,1]$
8	$[1,0,1]$	$[2,2,0]$	$[0,1,0]$	$[1,2,2]$	$[2,2,1]$	$[0,2,0]$
9	$[1,1,2]$	$[1,2,0]$	$[2,1,1]$	$[0,1,2]$	$[0,1,2]$	$[2,0,2]$
10	$[1,1,2]$	$[2,1,0]$	$[0,1,2]$	$[1,0,1]$	$[2,1,1]$	$[0,2,1]$
11	$[1,2,2]$	$[1,0,2]$	$[2,0,2]$	$[0,1,1]$	$[0,1,1]$	$[2,2,1]$
12	$[1,2,2]$	$[2,0,1]$	$[0,1,1]$	$[1,1,2]$	$[2,0,2]$	$[0,2,2]$
13	$[2,0,0]$	$[1,0,0]$	$[2,0,2]$	$[0,2,2]$	$[0,1,1]$	$[1,1,2]$
14	$[2,0,0]$	$[2,0,0]$	$[0,1,1]$	$[2,2,1]$	$[2,0,2]$	$[0,1,1]$
15	$[2,0,2]$	$[0,2,2]$	$[2,0,0]$	$[1,0,0]$	$[1,2,2]$	$[2,0,1]$
16	$[2,0,2]$	$[0,1,1]$	$[1,2,2]$	$[1,0,2]$	$[2,0,0]$	$[2,0,0]$
17	$[2,1,1]$	$[0,2,1]$	$[2,1,2]$	$[1,2,1]$	$[1,1,2]$	$[2,1,0]$
18	$[2,1,1]$	$[0,1,2]$	$[1,1,2]$	$[1,2,0]$	$[2,1,2]$	$[2,1,2]$
19	$[2,1,2]$	$[1,2,1]$	$[2,1,1]$	$[0,2,1]$	$[0,1,2]$	$[1,0,1]$
20	$[2,1,2]$	$[2,1,2]$	$[0,1,2]$	$[2,0,2]$	$[2,1,1]$	$[0,1,2]$
21	$[2,2,1]$	$[0,2,0]$	$[2,2,2]$	$[1,1,1]$	$[1,0,1]$	$[2,2,0]$
22	$[2,2,1]$	$[0,1,0]$	$[1,0,1]$	$[1,1,0]$	$[2,2,2]$	$[2,2,2]$
23	$[2,2,2]$	$[1,1,1]$	$[2,2,1]$	$[0,2,0]$	$[0,1,0]$	$[1,2,2]$
24	$[2,2,2]$	$[2,2,2]$	$[0,1,0]$	$[2,1,1]$	$[2,2,1]$	$[0,1,0]$

has the following five types

$$I: \quad p\mathcal{O}_K = \wp \qquad II: \quad p\mathcal{O}_K = \wp^3 \qquad III: \quad p\mathcal{O}_K = \wp_1\wp_2$$
$$IV: \quad p\mathcal{O}_K = \wp_1\wp_2^2 \quad V: \quad p\mathcal{O}_K = \wp_1\wp_2\wp_3$$

In type II, IV, V, all the primes over p have degree 1 residue field. In type I, the prime over p has degree 3 residue field. In type III, the degree of the residue fields of the two primes $\wp_1, \wp_2$ is 1 and 2 respectively. We choose the one with degree 1 residue field as $\wp$.

Suppose $x = (E, \pm P) \in Y_1(N)(K)$. Therefore by Lemma 3.5, E has good reduction at $\wp$. By Proposition 2.5, the reduction $\widetilde{P}$ of P is a k-rational point of order N in the elliptic curve $\widetilde{E}$ over $k = \mathcal{O}_K/\wp$.

The following lemma shows that $\widetilde{E}(k)$ can not have a point of order 24. This contradiction implies that $\mathbb{Z}/32\mathbb{Z}$ is not a subgroup of $E(K)_{tor}$. The calculations on the finite field $\mathbb{F}_{5^3}$ in this Lemma is done in Sage [31].

Lemma 4.2. *$\widetilde{E}(k)$ can not have a point of order 24.*

FIGURE 1. Atkin-Lehner involution on $\widetilde{X}_0(32)(\mathbb{F}_{3^3})$, $\omega_{32} = \omega$



FIGURE 2. Atkin-Lehner involution on $\widetilde{X}_0(32)(\mathbb{F}_{3^3})$, $\omega_{32} = \omega'$



Proof. For the modular curve $X_0(24)$, Fricke [6] calculated the defining equation and the explicit formula of j-invariant and the Atkin-Lehner involutions $\omega_4, \omega_9, \omega_{36}$ on this equation. These data can also be found in Furumoto-Hasegawa [7]. The equation is

$$y^2 = x^3 + 11x^2 + 36x + 36 \tag{4.2}$$

with the formula of the Atkin-Lehner involutions

$$\omega_{24}(P) = -P + (0, 6)$$
$$\omega_3(P) = P + (-3, 0)$$
$$\omega_8(P) = \omega_{24} \circ \omega_3^{-1}(P) = -P + (-4, 2)$$

and the formula of j-invariant

$$j = \frac{27\left(\frac{(x(x+6)/2)(2(x(x+6)/2)+9)^2}{27((x(x+6)/2)+4)}+1\right)\left(9\frac{(x(x+6)/2)(2(x(x+6)/2)+9)^2}{27((x(x+6)/2)+4)}+1\right)^3}{\frac{(x(x+6)/2)(2(x(x+6)/2)+9)^2}{27((x(x+6)/2)+4)}}$$

It can be calculated in Sage [31] that the set of rational torsion points of this equation is

$$\begin{aligned} X_0(24)(\mathbb{Q}) &= \{(\infty,\infty),(-4,2),(-2,0),(-4,-2),(-3,0),(0,6),(-6,0),(0,-6)\} \\ &\cong \mathbb{Z}/2\mathbb{Z}\times\mathbb{Z}/4\mathbb{Z} \end{aligned}$$

This is exactly all the rational points of this equation since $X_0(24)(\mathbb{Q}) \cong J_0(24)(\mathbb{Q})$ is finite.

The map $\pi : X_1(24) \longrightarrow X_0(24)$ factors through $X_\Delta(24)$, with $\Delta = \{\pm 1, \pm 11\}$, $g(X_\Delta(24)) = 1$ (See for example [11]). The cusps of $X_\Delta(24)$ over the 0 cusp of $X_0(24)$ are quadratic. This is because the four cusps of $X_1(24)$ over 0 are

$$\pm\begin{pmatrix}1\\0\end{pmatrix}, \pm\begin{pmatrix}5\\0\end{pmatrix}, \pm\begin{pmatrix}7\\0\end{pmatrix}, \pm\begin{pmatrix}11\\0\end{pmatrix}$$

They have two orbits under the action of Δ:

$$\left\{\pm\begin{pmatrix}1\\0\end{pmatrix}, \pm\begin{pmatrix}11\\0\end{pmatrix}\right\}, \quad \left\{\pm\begin{pmatrix}5\\0\end{pmatrix}, \pm\begin{pmatrix}7\\0\end{pmatrix}\right\}$$

So $X_\Delta(24)$ has two cusps $\pm\binom{1}{0}$, $\pm\binom{5}{0}$ over 0. They are conjugates since they have the same y in the representation $\pm\binom{x}{y}$.

The cusps of $X_\Delta(24)$ over the cusp $\pm\binom{1}{3}$ at $d = 3$ of $X_0(24)$ are quadratic. This is because the four cusps of $X_1(24)$ over $\pm\binom{1}{3}$ are

$$\pm\begin{pmatrix}1\\3\end{pmatrix}, \pm\begin{pmatrix}2\\3\end{pmatrix}, \pm\begin{pmatrix}1\\9\end{pmatrix}, \pm\begin{pmatrix}2\\9\end{pmatrix}$$

They have two orbits under the action of Δ:

$$\left\{\pm\begin{pmatrix}1\\3\end{pmatrix}, \pm\begin{pmatrix}2\\9\end{pmatrix}\right\}, \quad \left\{\pm\begin{pmatrix}2\\3\end{pmatrix}, \pm\begin{pmatrix}1\\9\end{pmatrix}\right\}$$

So $X_\Delta(24)$ has two cusps $\pm\binom{1}{3}$, $\pm\binom{2}{3}$ over $\pm\binom{1}{3}$. They are conjugates since they have the same y in the representation $\pm\binom{x}{y}$.

The cusps of $X_\Delta(24)$ over the cusp $\pm\binom{1}{8}$ at $d = 8$ of $X_0(24)$ are quadratic. This is because the four cusps of $X_1(24)$ over $\pm\binom{1}{8}$ are

$$\pm\begin{pmatrix}1\\8\end{pmatrix}, \pm\begin{pmatrix}3\\8\end{pmatrix}, \pm\begin{pmatrix}5\\8\end{pmatrix}, \pm\begin{pmatrix}7\\8\end{pmatrix}$$

They have two orbits under the action of Δ:

$$\left\{\pm\begin{pmatrix}1\\8\end{pmatrix}, \pm\begin{pmatrix}5\\8\end{pmatrix}\right\}, \quad \left\{\pm\begin{pmatrix}3\\8\end{pmatrix}, \pm\begin{pmatrix}7\\8\end{pmatrix}\right\}$$

So $X_\Delta(24)$ has two cusps $\pm\binom{1}{8}$, $\pm\binom{3}{8}$ over $\pm\binom{1}{8}$. They are conjugates since they have the same y in the representation $\pm\binom{x}{y}$.

Now $(0,6)=\omega_{24}(\infty)=0$, $(-3,0)=\omega_3(\infty)=\pm\binom{1}{3}$, $(-4,2)=\omega_8(\infty)=\pm\binom{1}{8}$ which means $(0,6),(-3,0)$ and $(-4,2)$ all come from quadratic cusps of $X_0(24)$ under $\pi_2: X_\Delta(24)\longrightarrow X_0(24)$.

If $k=\mathbb{F}_5$, then $\widetilde{E}(k)$ can not have a point of order 24 since $24>(1+\sqrt{5})^2$. If $k=\mathbb{F}_{5^3}$, let $[m,n,l]$ denote the element $ma^2+na+l\in\mathbb{F}_{5^3}=\mathbb{F}_5[a]$. Table 5 list the coordinates, j-invariant and φ of the points on the reduction of equation 4.2 modulo 5 which are rational over $\mathbb{F}_{5^3}$.

Table 5: Points on $\widetilde{X}_0(24)(\mathbb{F}_{5^3})$

X	Y_1	Y_2	j	$\varphi(X,Y_1)$	$\varphi(X,Y_2)$
∞	∞	∞	$-$		
0	4	1	$-$		
1	3	2	$-$		
2	0	0	$-$		
3	0	0	$-$		
4	0	0	$-$		
$[0,1,1]$	$[3,0,1]$	$[2,0,4]$	$[4,4,4]$	$(0,1)$	$(0,4)$
$[0,1,3]$	$[4,1,4]$	$[1,4,1]$	$[4,3,0]$	$(1,2)$	$(1,3)$
$[0,2,1]$	$[3,2,4]$	$[2,3,1]$	$[1,0,4]$	$(4,0)$	$(4,0)$
$[0,3,0]$	$[4,1,2]$	$[1,4,3]$	$[4,1,0]$	$(2,0)$	$(2,0)$
$[0,4,0]$	$[1,0,2]$	$[4,0,3]$	$[2,1,3]$	$(1,2)$	$(1,3)$
$[0,4,3]$	$[2,3,2]$	$[3,2,3]$	$[4,4,4]$	$(0,1)$	$(0,4)$
$[1,0,3]$	$[3,4,4]$	$[2,1,1]$	$[4,3,0]$	$(4,0)$	$(4,0)$
$[1,1,1]$	$[1,3,4]$	$[4,2,1]$	$[3,4,2]$	$(1,2)$	$(1,3)$
$[1,1,2]$	$[4,0,4]$	$[1,0,1]$	$[3,4,4]$	$(4,0)$	$(4,0)$
$[1,1,3]$	$[2,4,4]$	$[3,1,1]$	$[4,4,2]$	$(0,1)$	$(0,4)$
$[1,2,2]$	$[2,2,3]$	$[3,3,2]$	$[2,2,1]$	$(2,0)$	$(2,0)$
$[1,3,3]$	$[3,4,1]$	$[2,1,4]$	$[4,4,2]$	(∞,∞)	(∞,∞)
$[1,3,4]$	$[0,1,0]$	$[0,4,0]$	$[2,1,2]$	$(3,0)$	$(3,0)$
$[1,4,3]$	$[3,2,1]$	$[2,3,4]$	$[3,4,0]$	(∞,∞)	(∞,∞)
$[1,4,4]$	$[2,4,4]$	$[3,1,1]$	$[2,3,2]$	$(3,0)$	$(3,0)$
$[2,0,2]$	$[2,1,2]$	$[3,4,3]$	$[3,2,2]$	$(0,1)$	$(0,4)$
$[2,0,4]$	$[4,3,3]$	$[1,2,2]$	$[1,1,1]$	$(1,2)$	$(1,3)$
$[2,1,0]$	$[3,0,1]$	$[2,0,4]$	$[3,4,0]$	$(0,1)$	$(0,4)$
$[2,1,3]$	$[1,4,4]$	$[4,1,1]$	$[3,2,2]$	$(1,2)$	$(1,3)$
$[2,1,4]$	$[0,3,1]$	$[0,2,4]$	$[3,2,4]$	$(4,0)$	$(4,0)$
$[2,2,3]$	$[1,0,2]$	$[4,0,3]$	$[4,1,0]$	$(3,0)$	$(3,0)$
$[2,2,4]$	$[3,3,1]$	$[2,2,4]$	$[4,4,1]$	(∞,∞)	(∞,∞)
$[2,3,0]$	$[0,1,0]$	$[0,4,0]$	$[3,2,0]$	$(0,1)$	$(0,4)$
$[2,3,3]$	$[3,3,3]$	$[2,2,2]$	$[4,4,4]$	$(1,2)$	$(1,3)$
$[2,3,4]$	$[1,2,3]$	$[4,3,2]$	$[4,4,1]$	$(4,0)$	$(4,0)$
$[2,4,0]$	$[0,1,0]$	$[0,4,0]$	$[3,4,2]$	$(0,1)$	$(0,4)$
$[2,4,2]$	$[2,2,0]$	$[3,3,0]$	$[1,0,4]$	$(1,2)$	$(1,3)$
$[3,0,2]$	$[2,4,4]$	$[3,1,1]$	$[3,2,2]$	$(0,1)$	$(0,4)$
$[3,0,4]$	$[4,2,4]$	$[1,3,1]$	$[0,2,2]$	$(1,2)$	$(1,3)$
$[3,1,1]$	$[0,2,0]$	$[0,3,0]$	$[2,3,3]$	$(1,2)$	$(1,3)$

Table 5: continued

X	Y_1	Y_2	j	$\varphi(X,Y_1)$	$\varphi(X,Y_2)$
$[3,1,4]$	$[1,1,0]$	$[4,4,0]$	$[3,4,2]$	$(0,1)$	$(0,4)$
$[3,2,1]$	$[0,2,0]$	$[0,3,0]$	$[4,4,4]$	$(1,2)$	$(1,3)$
$[3,2,2]$	$[2,4,1]$	$[3,1,4]$	$[1,1,1]$	$(2,0)$	$(2,0)$
$[3,2,3]$	$[4,4,4]$	$[1,1,1]$	$[2,1,2]$	$(0,1)$	$(0,4)$
$[3,3,2]$	$[4,4,3]$	$[1,1,2]$	$[3,2,0]$	(∞,∞)	(∞,∞)
$[3,3,3]$	$[3,0,1]$	$[2,0,4]$	$[1,1,0]$	$(3,0)$	$(3,0)$
$[3,4,1]$	$[1,0,2]$	$[4,0,3]$	$[3,2,2]$	$(1,2)$	$(1,3)$
$[3,4,2]$	$[0,1,2]$	$[0,4,3]$	$[2,3,3]$	$(2,0)$	$(2,0)$
$[3,4,3]$	$[3,2,2]$	$[2,3,3]$	$[1,1,0]$	$(0,1)$	$(0,4)$
$[4,0,3]$	$[4,2,2]$	$[1,3,3]$	$[4,2,0]$	$(2,0)$	$(2,0)$
$[4,1,2]$	$[4,3,3]$	$[1,2,2]$	$[4,2,0]$	$(3,0)$	$(3,0)$
$[4,1,3]$	$[1,4,2]$	$[4,1,3]$	$[3,2,4]$	(∞,∞)	(∞,∞)
$[4,2,2]$	$[0,2,0]$	$[0,3,0]$	$[2,2,1]$	$(3,0)$	$(3,0)$
$[4,2,3]$	$[1,3,2]$	$[4,2,3]$	$[3,4,4]$	(∞,∞)	(∞,∞)
$[4,3,4]$	$[4,4,1]$	$[1,1,4]$	$[0,2,2]$	$(4,0)$	$(4,0)$
$[4,4,0]$	$[3,4,2]$	$[2,1,3]$	$[2,3,2]$	$(0,1)$	$(0,4)$
$[4,4,3]$	$[4,3,3]$	$[1,2,2]$	$[3,4,2]$	$(1,2)$	$(1,3)$
$[4,4,4]$	$[3,0,3]$	$[2,0,2]$	$[2,1,3]$	$(2,0)$	$(2,0)$

TABLE 6. Group structures of the elliptic curves over $\mathbb{F}_{5^3}$

j	$E_1(\mathbb{F}_{5^3})$	$E_2(\mathbb{F}_{5^3})$	j	$E_1(\mathbb{F}_{5^3})$	$E_2(\mathbb{F}_{5^3})$
$[0,2,2]$	$\mathbb{Z}/120$	$\mathbb{Z}/132$	$[1,0,4]$	$\mathbb{Z}/120$	$\mathbb{Z}/132$
$[1,1,0]$	$\mathbb{Z}/132$	$\mathbb{Z}/120$	$[1,1,1]$	$\mathbb{Z}/120$	$\mathbb{Z}/132$
$[2,1,2]$	$\mathbb{Z}/132$	$\mathbb{Z}/120$	$[2,1,3]$	$\mathbb{Z}/120$	$\mathbb{Z}/132$
$[2,2,1]$	$\mathbb{Z}/144$	$\mathbb{Z}/108$	$[2,3,2]$	$\mathbb{Z}/132$	$\mathbb{Z}/120$
$[2,3,3]$	$\mathbb{Z}/120$	$\mathbb{Z}/132$	$[3,2,0]$	$\mathbb{Z}/120$	$\mathbb{Z}/132$
$[3,2,2]$	$\mathbb{Z}/2\times\mathbb{Z}/72$	$\mathbb{Z}/2\times\mathbb{Z}/54$	$[3,2,4]$	$\mathbb{Z}/108$	$\mathbb{Z}/144$
$[3,4,0]$	$\mathbb{Z}/120$	$\mathbb{Z}/132$	$[3,4,2]$	$\mathbb{Z}/2\times\mathbb{Z}/72$	$\mathbb{Z}/2\times\mathbb{Z}/54$
$[3,4,4]$	$\mathbb{Z}/108$	$\mathbb{Z}/144$	$[4,1,0]$	$\mathbb{Z}/144$	$\mathbb{Z}/108$
$[4,2,0]$	$\mathbb{Z}/144$	$\mathbb{Z}/108$	$[4,3,0]$	$\mathbb{Z}/120$	$\mathbb{Z}/132$
$[4,4,1]$	$\mathbb{Z}/108$	$\mathbb{Z}/144$	$[4,4,2]$	$\mathbb{Z}/120$	$\mathbb{Z}/132$
$[4,4,4]$	$\mathbb{Z}/2\times\mathbb{Z}/72$	$\mathbb{Z}/2\times\mathbb{Z}/54$			

Table 6 shows the group structures of elliptic curves over $\mathbb{F}_{5^3}$ with the given j-invariants that appear in Table 5. It can be seen that all noncuspidal pionts on $\widetilde{X}_0(24)(\mathbb{F}_{5^3})$ may correspond to an elliptic curve with a point of order 24. By using the additive law on $X_0(24)$, $\omega_3(P)$, $\omega_8(P)$ and $\omega_{24}(P)$ are calculated as listed in Table 7.

Table 7: Atkin-Lehner involutions on $\widetilde{X}_0(24)(\mathbb{F}_{5^3})$

P	$X(P)$	$Y(P)$	$X(\omega_3(P))$	$Y(\omega_3(P))$	$X(\omega_8(P))$	$Y(\omega_8(P))$	$X(\omega_{24}(P))$	$Y(\omega_{24}(P))$
1	$(0,1,1)$	$(3,0,1)$	$(4,4,3)$	$(4,3,3)$	$(2,1,4)$	$(0,2,4)$	$(4,1,2)$	$(1,2,2)$
2	$(0,1,1)$	$(2,0,4)$	$(4,4,3)$	$(1,2,2)$	$(1,2,2)$	$(2,2,3)$	$(2,2,4)$	$(3,3,1)$
3	$(0,1,3)$	$(4,1,4)$	$(2,3,0)$	$(0,1,0)$	$(1,3,4)$	$(0,4,0)$	$(0,2,1)$	$(3,2,4)$
4	$(0,1,3)$	$(1,4,1)$	$(2,3,0)$	$(0,4,0)$	$(1,4,3)$	$(3,2,1)$	$(4,4,4)$	$(2,0,2)$
5	$(0,2,1)$	$(3,2,4)$	$(1,3,4)$	$(0,4,0)$	$(2,3,0)$	$(0,1,0)$	$(0,1,3)$	$(4,1,4)$
6	$(0,2,1)$	$(2,3,1)$	$(1,3,4)$	$(0,1,0)$	$(4,4,0)$	$(3,4,2)$	$(2,0,4)$	$(4,3,3)$
7	$(0,3,0)$	$(4,1,2)$	$(4,1,3)$	$(1,4,2)$	$(3,0,2)$	$(3,1,1)$	$(3,2,1)$	$(0,3,0)$
8	$(0,3,0)$	$(1,4,3)$	$(4,1,3)$	$(4,1,3)$	$(0,4,3)$	$(3,2,3)$	$(1,1,1)$	$(4,2,1)$
9	$(0,4,0)$	$(1,0,2)$	$(3,4,3)$	$(3,2,2)$	$(1,4,4)$	$(2,4,4)$	$(4,3,4)$	$(1,1,4)$
10	$(0,4,0)$	$(4,0,3)$	$(3,4,3)$	$(2,3,3)$	$(3,3,2)$	$(1,1,2)$	$(3,4,2)$	$(0,1,2)$
11	$(0,4,3)$	$(2,3,2)$	$(1,1,1)$	$(1,3,4)$	$(1,1,2)$	$(4,0,4)$	$(4,2,2)$	$(0,2,0)$
12	$(0,4,3)$	$(3,2,3)$	$(1,1,1)$	$(4,2,1)$	$(0,3,0)$	$(1,4,3)$	$(4,1,3)$	$(4,1,3)$
13	$(1,0,3)$	$(3,4,4)$	$(3,3,3)$	$(2,0,4)$	$(2,1,0)$	$(3,0,1)$	$(3,0,4)$	$(4,2,4)$
14	$(1,0,3)$	$(2,1,1)$	$(3,3,3)$	$(3,0,1)$	$(3,2,3)$	$(4,4,4)$	$(3,1,1)$	$(0,2,0)$
15	$(1,1,1)$	$(1,3,4)$	$(0,4,3)$	$(2,3,2)$	$(4,2,2)$	$(0,2,0)$	$(1,1,2)$	$(4,0,4)$
16	$(1,1,1)$	$(4,2,1)$	$(0,4,3)$	$(3,2,3)$	$(4,1,3)$	$(4,1,3)$	$(0,3,0)$	$(1,4,3)$
17	$(1,1,2)$	$(4,0,4)$	$(4,2,2)$	$(0,2,0)$	$(0,4,3)$	$(2,3,2)$	$(1,1,1)$	$(1,3,4)$
18	$(1,1,2)$	$(1,0,1)$	$(4,2,2)$	$(0,3,0)$	$(3,0,2)$	$(2,4,4)$	$(3,2,1)$	$(0,2,0)$
19	$(1,1,3)$	$(2,4,4)$	$(2,4,2)$	$(2,2,0)$	$(4,3,4)$	$(4,4,1)$	$(1,4,4)$	$(3,1,1)$
20	$(1,1,3)$	$(3,1,1)$	$(2,4,2)$	$(3,3,0)$	$(3,4,2)$	$(0,4,3)$	$(3,3,2)$	$(4,4,3)$
21	$(1,2,2)$	$(2,2,3)$	$(2,2,4)$	$(3,3,1)$	$(0,1,1)$	$(2,0,4)$	$(4,4,3)$	$(1,2,2)$
22	$(1,2,2)$	$(3,3,2)$	$(2,2,4)$	$(2,2,4)$	$(3,1,4)$	$(4,4,0)$	$(2,1,3)$	$(4,1,1)$
23	$(1,3,3)$	$(3,4,1)$	$(3,2,2)$	$(3,1,4)$	$(3,0,4)$	$(1,3,1)$	$(2,1,0)$	$(2,0,4)$
24	$(1,3,3)$	$(2,1,4)$	$(3,2,2)$	$(2,4,1)$	$(3,1,1)$	$(0,3,0)$	$(3,2,3)$	$(1,1,1)$
25	$(1,3,4)$	$(0,1,0)$	$(0,2,1)$	$(2,3,1)$	$(2,0,4)$	$(4,3,3)$	$(4,4,0)$	$(3,4,2)$
26	$(1,3,4)$	$(0,4,0)$	$(0,2,1)$	$(3,2,4)$	$(0,1,3)$	$(4,1,4)$	$(2,3,0)$	$(0,1,0)$
27	$(1,4,3)$	$(3,2,1)$	$(4,4,4)$	$(2,0,2)$	$(0,1,3)$	$(1,4,1)$	$(2,3,0)$	$(0,4,0)$
28	$(1,4,3)$	$(2,3,4)$	$(4,4,4)$	$(3,0,3)$	$(2,0,4)$	$(1,2,2)$	$(4,4,0)$	$(2,1,3)$
29	$(1,4,4)$	$(2,4,4)$	$(4,3,4)$	$(1,1,4)$	$(0,4,0)$	$(1,0,2)$	$(3,4,3)$	$(3,2,2)$
30	$(1,4,4)$	$(3,1,1)$	$(4,3,4)$	$(4,4,1)$	$(2,4,2)$	$(2,2,0)$	$(1,1,3)$	$(2,4,4)$
31	$(2,0,2)$	$(2,1,2)$	$(2,3,3)$	$(3,3,3)$	$(2,3,4)$	$(1,2,3)$	$(2,2,3)$	$(1,0,2)$
32	$(2,0,2)$	$(3,4,3)$	$(2,3,3)$	$(2,2,2)$	$(4,0,3)$	$(1,3,3)$	$(4,2,3)$	$(4,2,3)$
33	$(2,0,4)$	$(4,3,3)$	$(4,4,0)$	$(3,4,2)$	$(1,3,4)$	$(0,1,0)$	$(0,2,1)$	$(2,3,1)$
34	$(2,0,4)$	$(1,2,2)$	$(4,4,0)$	$(2,1,3)$	$(1,4,3)$	$(2,3,4)$	$(4,4,4)$	$(3,0,3)$
35	$(2,1,0)$	$(3,0,1)$	$(3,0,4)$	$(4,2,4)$	$(1,0,3)$	$(3,4,4)$	$(3,3,3)$	$(2,0,4)$
36	$(2,1,0)$	$(2,0,4)$	$(3,0,4)$	$(1,3,1)$	$(3,2,2)$	$(3,1,4)$	$(1,3,3)$	$(3,4,1)$
37	$(2,1,3)$	$(1,4,4)$	$(3,1,4)$	$(1,1,0)$	$(4,1,2)$	$(4,3,3)$	$(2,1,4)$	$(0,3,1)$
38	$(2,1,3)$	$(4,1,1)$	$(3,1,4)$	$(4,4,0)$	$(2,2,4)$	$(2,2,4)$	$(1,2,2)$	$(3,3,2)$
39	$(2,1,4)$	$(0,3,1)$	$(4,1,2)$	$(4,3,3)$	$(3,1,4)$	$(1,1,0)$	$(2,1,3)$	$(1,4,4)$
40	$(2,1,4)$	$(0,2,4)$	$(4,1,2)$	$(1,2,2)$	$(0,1,1)$	$(3,0,1)$	$(4,4,3)$	$(4,3,3)$
41	$(2,2,3)$	$(1,0,2)$	$(2,3,4)$	$(1,2,3)$	$(2,3,3)$	$(3,3,3)$	$(2,0,2)$	$(2,1,2)$
42	$(2,2,3)$	$(4,0,3)$	$(2,3,4)$	$(4,3,2)$	$(3,4,1)$	$(1,0,2)$	$(2,4,0)$	$(0,1,0)$
43	$(2,2,4)$	$(3,3,1)$	$(1,2,2)$	$(2,2,3)$	$(4,4,3)$	$(1,2,2)$	$(0,1,1)$	$(2,0,4)$

Table 7: continued

P	$X(P)$	$Y(P)$	$X(\omega_3(P))$	$Y(\omega_3(P))$	$X(\omega_8(P))$	$Y(\omega_8(P))$	$X(\omega_{24}(P))$	$Y(\omega_{24}(P))$
44	$(2,2,4)$	$(2,2,4)$	$(1,2,2)$	$(3,3,2)$	$(2,1,3)$	$(4,1,1)$	$(3,1,4)$	$(4,4,0)$
45	$(2,3,0)$	$(0,1,0)$	$(0,1,3)$	$(4,1,4)$	$(0,2,1)$	$(3,2,4)$	$(1,3,4)$	$(0,4,0)$
46	$(2,3,0)$	$(0,4,0)$	$(0,1,3)$	$(1,4,1)$	$(4,4,4)$	$(2,0,2)$	$(1,4,3)$	$(3,2,1)$
47	$(2,3,3)$	$(3,3,3)$	$(2,0,2)$	$(2,1,2)$	$(2,2,3)$	$(1,0,2)$	$(2,3,4)$	$(1,2,3)$
48	$(2,3,3)$	$(2,2,2)$	$(2,0,2)$	$(3,4,3)$	$(4,2,3)$	$(4,2,3)$	$(4,0,3)$	$(1,3,3)$
49	$(2,3,4)$	$(1,2,3)$	$(2,2,3)$	$(1,0,2)$	$(2,0,2)$	$(2,1,2)$	$(2,3,3)$	$(3,3,3)$
50	$(2,3,4)$	$(4,3,2)$	$(2,2,3)$	$(4,0,3)$	$(2,4,0)$	$(0,1,0)$	$(3,4,1)$	$(1,0,2)$
51	$(2,4,0)$	$(0,1,0)$	$(3,4,1)$	$(1,0,2)$	$(2,3,4)$	$(4,3,2)$	$(2,2,3)$	$(4,0,3)$
52	$(2,4,0)$	$(0,4,0)$	$(3,4,1)$	$(4,0,3)$	$(4,0,3)$	$(4,2,2)$	$(4,2,3)$	$(1,3,2)$
53	$(2,4,2)$	$(2,2,0)$	$(1,1,3)$	$(2,4,4)$	$(1,4,4)$	$(3,1,1)$	$(4,3,4)$	$(4,4,1)$
54	$(2,4,2)$	$(3,3,0)$	$(1,1,3)$	$(3,1,1)$	$(3,3,2)$	$(4,4,3)$	$(3,4,2)$	$(0,4,3)$
55	$(3,0,2)$	$(2,4,4)$	$(3,2,1)$	$(0,2,0)$	$(1,1,2)$	$(1,0,1)$	$(4,2,2)$	$(0,3,0)$
56	$(3,0,2)$	$(3,1,1)$	$(3,2,1)$	$(0,3,0)$	$(0,3,0)$	$(4,1,2)$	$(4,1,3)$	$(1,4,2)$
57	$(3,0,4)$	$(4,2,4)$	$(2,1,0)$	$(3,0,1)$	$(3,3,3)$	$(2,0,4)$	$(1,0,3)$	$(3,4,4)$
58	$(3,0,4)$	$(1,3,1)$	$(2,1,0)$	$(2,0,4)$	$(1,3,3)$	$(3,4,1)$	$(3,2,2)$	$(3,1,4)$
59	$(3,1,1)$	$(0,2,0)$	$(3,2,3)$	$(4,4,4)$	$(3,3,3)$	$(3,0,1)$	$(1,0,3)$	$(2,1,1)$
60	$(3,1,1)$	$(0,3,0)$	$(3,2,3)$	$(1,1,1)$	$(1,3,3)$	$(2,1,4)$	$(3,2,2)$	$(2,4,1)$
61	$(3,1,4)$	$(1,1,0)$	$(2,1,3)$	$(1,4,4)$	$(2,1,4)$	$(0,3,1)$	$(4,1,2)$	$(4,3,3)$
62	$(3,1,4)$	$(4,4,0)$	$(2,1,3)$	$(4,1,1)$	$(1,2,2)$	$(3,3,2)$	$(2,2,4)$	$(2,2,4)$
63	$(3,2,1)$	$(0,2,0)$	$(3,0,2)$	$(2,4,4)$	$(4,2,2)$	$(0,3,0)$	$(1,1,2)$	$(1,0,1)$
64	$(3,2,1)$	$(0,3,0)$	$(3,0,2)$	$(3,1,1)$	$(4,1,3)$	$(1,4,2)$	$(0,3,0)$	$(4,1,2)$
65	$(3,2,2)$	$(2,4,1)$	$(1,3,3)$	$(2,1,4)$	$(3,2,3)$	$(1,1,1)$	$(3,1,1)$	$(0,3,0)$
66	$(3,2,2)$	$(3,1,4)$	$(1,3,3)$	$(3,4,1)$	$(2,1,0)$	$(2,0,4)$	$(3,0,4)$	$(1,3,1)$
67	$(3,2,3)$	$(4,4,4)$	$(3,1,1)$	$(0,2,0)$	$(1,0,3)$	$(2,1,1)$	$(3,3,3)$	$(3,0,1)$
68	$(3,2,3)$	$(1,1,1)$	$(3,1,1)$	$(0,3,0)$	$(3,2,2)$	$(2,4,1)$	$(1,3,3)$	$(2,1,4)$
69	$(3,3,2)$	$(4,4,3)$	$(3,4,2)$	$(0,4,3)$	$(2,4,2)$	$(3,3,0)$	$(1,1,3)$	$(3,1,1)$
70	$(3,3,2)$	$(1,1,2)$	$(3,4,2)$	$(0,1,2)$	$(0,4,0)$	$(4,0,3)$	$(3,4,3)$	$(2,3,3)$
71	$(3,3,3)$	$(3,0,1)$	$(1,0,3)$	$(2,1,1)$	$(3,1,1)$	$(0,2,0)$	$(3,2,3)$	$(4,4,4)$
72	$(3,3,3)$	$(2,0,4)$	$(1,0,3)$	$(3,4,4)$	$(3,0,4)$	$(4,2,4)$	$(2,1,0)$	$(3,0,1)$
73	$(3,4,1)$	$(1,0,2)$	$(2,4,0)$	$(0,1,0)$	$(2,2,3)$	$(4,0,3)$	$(2,3,4)$	$(4,3,2)$
74	$(3,4,1)$	$(4,0,3)$	$(2,4,0)$	$(0,4,0)$	$(4,2,3)$	$(1,3,2)$	$(4,0,3)$	$(4,2,2)$
75	$(3,4,2)$	$(0,1,2)$	$(3,3,2)$	$(1,1,2)$	$(3,4,3)$	$(2,3,3)$	$(0,4,0)$	$(4,0,3)$
76	$(3,4,2)$	$(0,4,3)$	$(3,3,2)$	$(4,4,3)$	$(1,1,3)$	$(3,1,1)$	$(2,4,2)$	$(3,3,0)$
77	$(3,4,3)$	$(3,2,2)$	$(0,4,0)$	$(1,0,2)$	$(4,3,4)$	$(1,1,4)$	$(1,4,4)$	$(2,4,4)$
78	$(3,4,3)$	$(2,3,3)$	$(0,4,0)$	$(4,0,3)$	$(3,4,2)$	$(0,1,2)$	$(3,3,2)$	$(1,1,2)$
79	$(4,0,3)$	$(4,2,2)$	$(4,2,3)$	$(1,3,2)$	$(2,4,0)$	$(0,4,0)$	$(3,4,1)$	$(4,0,3)$
80	$(4,0,3)$	$(1,3,3)$	$(4,2,3)$	$(4,2,3)$	$(2,0,2)$	$(3,4,3)$	$(2,3,3)$	$(2,2,2)$
81	$(4,1,2)$	$(4,3,3)$	$(2,1,4)$	$(0,3,1)$	$(2,1,3)$	$(1,4,4)$	$(3,1,4)$	$(1,1,0)$
82	$(4,1,2)$	$(1,2,2)$	$(2,1,4)$	$(0,2,4)$	$(4,4,3)$	$(4,3,3)$	$(0,1,1)$	$(3,0,1)$
83	$(4,1,3)$	$(1,4,2)$	$(0,3,0)$	$(4,1,2)$	$(3,2,1)$	$(0,3,0)$	$(3,0,2)$	$(3,1,1)$
84	$(4,1,3)$	$(4,1,3)$	$(0,3,0)$	$(1,4,3)$	$(1,1,1)$	$(4,2,1)$	$(0,4,3)$	$(3,2,3)$
85	$(4,2,2)$	$(0,2,0)$	$(1,1,2)$	$(4,0,4)$	$(1,1,1)$	$(1,3,4)$	$(0,4,3)$	$(2,3,2)$
86	$(4,2,2)$	$(0,3,0)$	$(1,1,2)$	$(1,0,1)$	$(3,2,1)$	$(0,2,0)$	$(3,0,2)$	$(2,4,4)$
87	$(4,2,3)$	$(1,3,2)$	$(4,0,3)$	$(4,2,2)$	$(3,4,1)$	$(4,0,3)$	$(2,4,0)$	$(0,4,0)$

Table 7: continued

P	$X(P)$	$Y(P)$	$X(\omega_3(P))$	$Y(\omega_3(P))$	$X(\omega_8(P))$	$Y(\omega_8(P))$	$X(\omega_{24}(P))$	$Y(\omega_{24}(P))$
88	$(4,2,3)$	$(4,2,3)$	$(4,0,3)$	$(1,3,3)$	$(2,3,3)$	$(2,2,2)$	$(2,0,2)$	$(3,4,3)$
89	$(4,3,4)$	$(4,4,1)$	$(1,4,4)$	$(3,1,1)$	$(1,1,3)$	$(2,4,4)$	$(2,4,2)$	$(2,2,0)$
90	$(4,3,4)$	$(1,1,4)$	$(1,4,4)$	$(2,4,4)$	$(3,4,3)$	$(3,2,2)$	$(0,4,0)$	$(1,0,2)$
91	$(4,4,0)$	$(3,4,2)$	$(2,0,4)$	$(4,3,3)$	$(0,2,1)$	$(2,3,1)$	$(1,3,4)$	$(0,1,0)$
92	$(4,4,0)$	$(2,1,3)$	$(2,0,4)$	$(1,2,2)$	$(4,4,4)$	$(3,0,3)$	$(1,4,3)$	$(2,3,4)$
93	$(4,4,3)$	$(4,3,3)$	$(0,1,1)$	$(3,0,1)$	$(4,1,2)$	$(1,2,2)$	$(2,1,4)$	$(0,2,4)$
94	$(4,4,3)$	$(1,2,2)$	$(0,1,1)$	$(2,0,4)$	$(2,2,4)$	$(3,3,1)$	$(1,2,2)$	$(2,2,3)$
95	$(4,4,4)$	$(3,0,3)$	$(1,4,3)$	$(2,3,4)$	$(4,4,0)$	$(2,1,3)$	$(2,0,4)$	$(1,2,2)$
96	$(4,4,4)$	$(2,0,2)$	$(1,4,3)$	$(3,2,1)$	$(2,3,0)$	$(0,4,0)$	$(0,1,3)$	$(1,4,1)$

The information in Table 7 can be represented in Figure 3. The 96-vertex graph has 24 connected components, which are all complete graphs of order 4. The black dot points are those such that

$$\varphi(P) \in \{(-3,0),(-4,2),(0,6)\}$$

while the white dot points are those such that

$$\varphi(P) \notin \{(-3,0),(-4,2),(0,6)\}$$

The horizontal lines represent ω_3. The vertical lines represent ω_8. And the diagonal lines represent ω_{24}. It is clear that each white dot point is connected to at least one black dot point, which is not allowed by Lemma 3.8.

□

Acknowledgements. It is a pleasure to thank my PhD advisor Sheldon Kamienny for introducing this research topic and for providing many valuable ideas and insightful comments throughout the research. I also wish to thank Andrew Sutherland, Andreas Schweizer and Maarten Derickx for pointing out several errors in an earlier version of this paper.

REFERENCES

[1] Bosch, S.; Lütkebohmert, W.; Raynaud, M.: *Néron models.* Ergebnisse der Mathematik und ihrer Grenzgebiete (3), 21. Springer-Verlag, Berlin (1990)

[2] Cassels, J. W. S.: *A note on the division values of* $\wp(u)$. Proc. Cambridge Philos. Soc. 45, 167-172 (1949)

[3] Diamond, F., Shurman, J.: *A first course in modular forms.* Graduate Texts in Mathematics, 228. Springer-Verlag, New York (2005)

[4] Drinfeld, V. G.: *Two theorems on modular curves.* (Russian) Funkcional. Anal. i Priložen. 7, no. 2, 83-84 (1973)

FIGURE 3. Atkin-Lehner involutions on $\widetilde{X}_0(24)(\mathbb{F}_{5^3})$



[5] Frey, G.: *Curves with infinitely many points of fixed degree.* Israel J. Math. 85, no. 1-3, 79-83 (1994)

[6] Fricke, R.: *Die elliptischen Funktionen und ihre Anwendungen. Erster Teil. Die funktionentheoretischen und analytischen Grundlagen. (German)*©2012. Reprint of the 1916 original. Springer, Heidelberg (2011)

[7] Furumoto, M.; Hasegawa, Y.: *Hyperelliptic quotients of modular curves $X_0(N)$.* Tokyo J. Math. 22, no. 1, 105-125 (1999)

[8] Igusa, J.: *Kroneckerian model of fields of elliptic modular functions.* Amer. J. Math. 81, 561-577 (1959)

[9] Ishii, N., Momose, F.: *Hyperelliptic modular curves.* Tsukuba J. Math. 15, no. 2, 413-423 (1991)

[10] Jeon, D.; Kim, C. H.: *Bielliptic modular curves $X_1(N)$.* Acta Arith. 112, no. 1, 75-86 (2004)

[11] Jeon, D.; Kim, C. H.: *On the arithmetic of certain modular curves.* Acta Arith. 130, no. 2, 181-193 (2007)

[12] Jeon, D., Kim, C. H., Schweizer, A.: *On the torsion of elliptic curves over cubic number fields.* Acta Arith. 113, no. 3, 291-301 (2004)

[13] Jeong, E. K.: *Isomorphism classes of elliptic curves over finite feilds with characteristic 3.* J. Chungcheong Math. Soc. 22, no. 3, 299-307 (2009)

[14] Kamienny, S.: *Torsion points on elliptic curves and q-coefficients of modular forms.* Invent. Math. 109, no. 2, 221-229 (1992)

[15] Kato, K.: *p-adic Hodge theory and values of zeta functions of modular forms.* Astérisque, (295):ix, 117-290. Cohomologies p-adiques et applications arithmétiques. III (2004)

[16] Katz, N. M.: *Galois properties of torsion points on abelian varieties.* Invent. Math. 62 , no. 3, 481-502 (1981)

[17] Kenku, M. A., Momose, F.: *Torsion points on elliptic curves defined over quadratic fields.* Nagoya Math. J. 109, 125-149 (1988)

[18] Kim, C. H.; Koo, J. K.: *The normalizer of $\Gamma_1(N)$ in $PSL_2(\mathbb{R})$.* Comm. Algebra 28, no. 11, 5303-5310 (2000)

[19] Kodaira, K. *On compact analytic surfaces: II.* Ann. of Math. (2) 77, 563-626 (1963)

[20] Kubert, D. S.: *Universal bounds on the torsion of elliptic curves.* Proc. London Math. Soc. (3) 33, no. 2, 193-237 (1976)

[21] Magma: http://magma.maths.usyd.edu.au/magma/

[22] Manin, Y. I.: *Parabolic points and zeta functions of modular curves.* (Russian) Izv. Akad. Nauk SSSR Ser. Mat. 36, 19-66 (1972)

[23] Mazur, B.: *Modular curves and the Eisenstein ideal.* Inst. Hautes Études Sci. Publ. Math. No. 47 (1977), 33-186 (1978)

[24] Najman, F.: *Torsion of rational elliptic curves over cubic fields and sporadic points on $X_1(n)$.* Math. Res. Lett. 23, no. 1, 245-272 (2016)

[25] Néron, A.: *Modèles minimaux des variétés abéliennes sur les corps locaux et globaux.* (French) Inst. Hautes Études Sci. Publ.Math. No. 21, 1-128 (1964)

[26] Ogg, A. P.: *Rational points on certain elliptic modular curves.* Analytic number theory (Proc. Sympos. Pure Math., Vol XXIV, St. Louis Univ., St. Louis, Mo., 1972), pp. 221-231. Amer. Math. Soc., Providence, R.I. (1973)

[27] Ogg, A. P.: *Hyperelliptic modular curves.* Bull. Soc. Math. France 102, 449-462 (1974)

[28] Ogg, A. P.: *Diophantine equations and modular forms.* Bull. Amer. Math. Soc. 81, 14-27 (1975)

[29] Parent, P.: *Torsion des courbes elliptiques sur les corps cubiques. (French)* Ann. Inst. Fourier (Grenoble) 50, no. 3, 723-749 (2000)

[30] Parent, P.: *No 17-torsion on elliptic curves over cubic number fields.* J. Théor. Nombres Bordeaux 15, no. 3, 831-838 (2003)

[31] Sage: http://www.sagemath.org/

[32] Shimura, G.: *Introduction to the arithmetic theory of automorphic functions.* Kanô Memorial Lectures, No. 1. Publications of the Mathematical Society of Japan, No. 11. Iwanami Shoten, Publishers, Tokyo; Princeton University Press, Princeton, N.J. (1971)

[33] Silverman, J. H.: *The arithmetic of elliptic curves.* Second edition. Graduate Texts in Mathematics, 106. Springer, Dordrecht (2009)

[34] Silverman, J. H.: *Advanced topics in the arithmetic of elliptic curves.* Graduate Texts in Mathematics, 151. Springer-Verlag, New York (1994)

[35] Waterhouse, W. C.: *Abelian varieties over finite fields.* Ann. Sci. École Norm. Sup. (4) 2 521-560 (1969)

DEPARTMENT OF MATHEMATICAL SCIENCES, TSINGHUA UNIVERSITY, BEIJING, 100084, CHINA
E-mail address: blandye@gmail.com